



Janet Napolitano, Governor  
Stephen A. Owens, ADEQ Director

# 2006 AIR QUALITY ANNUAL REPORT

(A.R.S. 49-424.10)

# ***Air Quality Report***

## ***A.R.S. §49-424.10***

---

### **Acknowledgments**

---

Numerous agencies, companies, individuals and organizations have collected the ambient air quality monitoring data presented in this report. The Arizona Department of Environmental Quality (ADEQ) publishes data from these various sources to provide as complete a picture as possible of air quality conditions throughout Arizona and gratefully acknowledges the efforts of all involved. Generally, ambient data presented in this report are collected, processed and reported following U.S. Environmental Protection Agency (EPA) policies and procedures. Air quality data that ADEQ staff and contract operators collect have also received internal and external quality control and assurance checks. Data provided by other sources have been checked by the responsible organization but not by ADEQ.

Private individuals and companies under contract to ADEQ provided invaluable field sampler operation and data processing services in support of monitoring activities during 2005. Their efforts are appreciated as they maneuvered on rooftops and metal towers to operate monitoring equipment in uncomfortable weather conditions, or review instrument performance and ambient monitoring data for technical accuracy. Field staff from other public agencies also operated numerous ambient monitoring sites in Arizona, providing spatial resolution and temporal coverage of air quality conditions statewide. ADEQ recognizes the efforts of these other monitoring and reporting agencies, and appreciates the opportunity to publish their data. Several industrial facilities collected and reported ambient air quality data to ADEQ, usually to satisfy permit requirements; their efforts are also acknowledged. Finally, ADEQ staff work daily installing, calibrating, maintaining, conducting quality control checks, collecting, processing, performing quality assurance tests and reporting data from a wide variety of ambient air monitoring instruments. ADEQ management wishes to thank these staff members for their dedication to maintaining and improving the quality of our program.

This report was prepared by ADEQ's Air Quality Assessment Section, which can be contacted at 1110 W. Washington St., Phoenix, AZ 85007, (602) 771-2274 or, toll free in Arizona at (800) 234-5677, then enter 771-2274. Our Web site is located at <http://www.azdeq.gov/>.

## Introduction

---

This report presents the results of air quality monitoring conducted throughout Arizona in the 2005 calendar year. Data from more than 100 monitoring sites are included in this report. In addition to the ADEQ monitoring network, air quality agencies in Maricopa, Pima and Pinal counties also operated networks, as did several industrial facilities. Their data are summarized in this report. Many of the sites have multiple instruments measuring a variety of gaseous, particulate and visibility parameters. The majority of the air quality measurements are for criteria pollutants (ozone, particulate matter, sulfur dioxide, carbon monoxide, nitrogen dioxide and lead) for which EPA has established National Ambient Air Quality Standards (NAAQS). Visibility-related measurements are included from a statewide network of operators.

The report on ambient air quality monitoring networks, which begins on Page 3, discusses the purpose, measurement methods and the specific scale of geographic resolution of each network of various air monitoring networks in Arizona.

Beginning on Page 14, the monitoring data report summarizes the monitoring data and shows the compliance status for criteria pollutants. It consists of three sections: measurement of traditional criteria pollutants, compliance status of the criteria pollutants, and visibility characterization. The text describes how the measurements are made and how they relate to compliance with the NAAQS.

The report on special projects, which begins on Page 65, summarizes activities from special monitoring projects undertaken in the last few years which have continued into 2005. Some of the projects presented in this report are the expanding Class I visibility monitoring network for larger national parks and wilderness areas, a new and expanding effort to characterize ozone precursors, and an intensive ambient monitoring and risk assessment project beginning in the Yuma area.

Air quality trends are reported beginning on Page 70. Air quality trends at most of the long-term monitors reveal improved air quality. Concentrations of carbon monoxide, lead and sulfur dioxide have improved dramatically since measurements began in the 1970s, and all monitors for these pollutants have shown compliance with health standards in recent years. Particulate matter (PM<sub>10</sub>) concentrations have also improved in rural and industrial areas where controls have been implemented, while less dramatic improvements have occurred in the neighborhoods of Phoenix and Tucson. Ozone concentrations have been fairly steady in Tucson and Yuma but have decreased since 1997 in Phoenix. On May 30, 2001, Maricopa County reached attainment for the 1-hour ozone standard. Effective June 15, 2004, the Phoenix area was designated nonattainment for the 8-hour ozone standard. Shorter periods of record for visibility in the urban and national parks and wilderness areas make trend assessments less definitive, but trend assessments are shown for the two urban areas.

## Ambient Air Quality Monitoring Networks

---

The federal Clean Air Act of 1970 required EPA to assist states and localities in establishing ambient air quality monitoring networks to characterize human health exposure and public welfare effects of criteria pollutants. The 1977 federal Clean Air Act amendments required each state to implement a visibility monitoring network to cover specified national parks and wilderness areas. The Phoenix and Tucson

metropolitan areas also have year-round visibility monitoring networks to assess urban haze. All of these networks are composed of individual monitoring sites; they are operated to collect ambient air quality data to ensure that Arizona citizens are able to know local air quality conditions and help ADEQ and local air quality control agencies identify the causes of polluted air.



**Figure 1 – Greer visibility monitoring site, located at 8,255 feet elevation in the Mt. Baldy Wilderness Area.**

### Criteria Pollutant Monitoring Networks

Ambient monitoring networks for air quality are established to sample pollution in a variety of representative settings, to assess the health and welfare effects, and to assist in determining air pollution sources. These networks cover both urban and rural areas of the state. Sampling networks are designed to satisfy monitoring objectives and measurement scales defined in Tables 1 and 2. Networks operated to monitor the nature and causes of visibility impairment use some of the same sampling methods and are described in more detail later in this section.

The criteria pollutants are presently defined as carbon monoxide (CO), nitrogen dioxide (NO<sub>2</sub>), sulfur dioxide (SO<sub>2</sub>), ozone (O<sub>3</sub>), suspended particulate matter (PM), and total particulate lead (Pb). These pollutants are monitored with federal reference or equivalent methods that EPA has certified. EPA defined particulate matter monitoring in 1987 to measure particles less than or equal to 10 microns in aerodynamic diameter (PM<sub>10</sub>), and again in 1997 to measure both PM<sub>10</sub> and, separately, particles less than or equal to 2.5 microns in aerodynamic diameter (PM<sub>2.5</sub>).

For each criteria pollutant, EPA specifies monitoring objectives that define the parameters by which health exposure and public welfare are assessed and the measurement scale classifications that describe the influence of atmospheric movement at a given location.

The types and scales of monitoring sites described above are combined into networks, which a number of government agencies and regulated companies operate. These networks are composed of one or more monitoring sites whose data are compared to the NAAQS and statistically analyzed in various ways. The agency or company operating a monitoring network also tracks data recovery, quality control and quality assurance parameters for the instruments operated at their various sites. The agency or company also often measures meteorological variables at the monitoring site.

Number	Definition
1	Determine highest concentrations expected to occur in the area covered by the network
2	Determine representative concentrations in areas of high population density
3	Determine the impact on ambient pollution levels of significant sources or source categories
4	Determine general background concentration levels
5	Determine the extent of regional pollutant transport among populated areas and in support of secondary standards
6	Determine the welfare-related effects in more rural and remote areas (such as visibility impairment and vegetation effects)

Measurement Scale <i>represents concentrations in air volumes within areas defined below</i>	Criteria Pollutant					
	Carbon Monoxide (CO)	Nitrogen Dioxide (NO <sub>2</sub> )	Ozone (O <sub>3</sub> )	Sulfur Dioxide (SO <sub>2</sub> )	Particulate Matter (PM <sub>10</sub> , PM <sub>2.5</sub> )	Lead (Pb)
Micro (0 to 100 m)	X				X	X
Middle (~100 to 500 m)	X	X	X	X	X	X
Neighborhood (~0.5 to 4 km)	X	X	X	X	X	X
Urban (~4 to 50 km)		X	X	X	X	X
Regional (~10 to 100s of km)			X	X	X	X

Some of the agencies do special continuous monitoring for the optical characteristics of the atmosphere and manual sampling of ozone-forming compounds and other hazardous air pollutants. Maricopa, Pima and Pinal counties operate networks

primarily to monitor urban air pollution. In contrast, the industrial networks are operated to determine the effects of their emissions on local air quality. The National Park Service's network tracks conditions in and around national parks and monuments. The state network monitors a wide variety of pollutant and atmospheric characteristics, including urban, industrial, rural and background surveillance.

The monitoring networks and their characteristics are shown in Table 3. A list of individual sites and monitoring parameters, based on the best available information at the time of publication, is presented in Appendix 1.

<i>Table 3. Monitoring Networks Operating in Arizona</i>				
Network Operator	Geographic Area Monitored	Monitoring Objective*	Measurement Scale(s)**	Pollutant(s) Monitored
Arizona Dept. of Environmental Quality	Statewide	1, 2, 3, 4, 5, 6	Micro, Middle, Neighborhood, Urban, Regional	SO <sub>2</sub> , O <sub>3</sub> , NO <sub>2</sub> , CO, PM <sub>10</sub> , PM <sub>2.5</sub>
Arizona Portland Cement Company	Rillito	1, 3	Neighborhood	PM <sub>10</sub>
ASARCO LLC.	Hayden	1, 2, 3	Middle, Neighborhood	SO <sub>2</sub>
Maricopa County Environmental Services Dept.	Phoenix urban area, Maricopa County	1, 2, 3, 4, 5, 6	Micro, Middle, Neighborhood, Urban, Regional	SO <sub>2</sub> , O <sub>3</sub> , NO <sub>2</sub> , CO, PM <sub>10</sub> , PM <sub>2.5</sub>
National Park Service	National parks and monuments	3, 4, 5, 6	Urban, Regional	SO <sub>2</sub> , O <sub>3</sub> , NO <sub>2</sub> , PM <sub>10</sub> , PM <sub>2.5</sub>
Phelps Dodge Miami Inc. (PDMI)	Miami	1, 2, 3	Neighborhood	SO <sub>2</sub> , PM <sub>10</sub> , PM <sub>2.5</sub>
Phoenix Cement Company	Clarkdale	1, 3	Neighborhood	PM <sub>10</sub>
Pima County Dept. of Environmental Quality	Tucson urban area, Pima County	1, 2, 3, 4, 5, 6	Micro, Middle, Neighborhood, Urban, Regional	SO <sub>2</sub> , O <sub>3</sub> , NO <sub>2</sub> , CO, PM <sub>10</sub> , PM <sub>2.5</sub>
Pinal County Air Quality Control District	Pinal County, Phoenix urban area	1, 2, 3, 4, 5	Middle, Neighborhood, Urban, Regional	O <sub>3</sub> , CO, PM <sub>10</sub> , PM <sub>2.5</sub>


<i>Table 3. Monitoring Networks Operating in Arizona</i>				
Network Operator	Geographic Area Monitored	Monitoring Objective*	Measurement Scale(s)**	Pollutant(s) Monitored
Praxair, Inc.	Kingman	1, 3	Middle	PM <sub>10</sub>
Salt River Project	Page	1, 3	Urban, Regional	NO <sub>2</sub> , O <sub>3</sub> , SO <sub>2</sub> , PM <sub>10</sub> , PM <sub>2.5</sub>
Tucson Electric Power Company	Tucson and Springerville	1, 2, 3	Middle, Regional	SO <sub>2</sub> , NO <sub>2</sub> , PM <sub>10</sub> , PM <sub>2.5</sub>

\*See Table 1 for a list of monitoring objectives

\*\*See Table 2 for a definition of measurement scales

### Visibility Monitoring Networks in National Parks and Wilderness Areas

The intent of the Class I visibility monitoring program is to characterize long-term trends as completely as possible using ambient visibility measurements within constraints of an area's size, terrain or logistics for each of the 12 federally protected Class I areas in Arizona (see Figure 2 and Appendix 4). The objectives of the visibility monitoring network are to track short-term and long-term trends in Arizona Class I areas, to assist in identifying any visibility impairment caused by existing major industrial sources, and to provide monitoring data if necessary for new or major modifications of major industrial sources.



**Figure 2 - Visibility Monitoring Sites Statewide**

Arizona continues to participate in the Interagency Monitoring of Protected Visual Environments (IMPROVE) Program as part of the overall national visibility monitoring effort. IMPROVE is a cooperative measurement effort between EPA, federal land management agencies and state air agencies. The objectives of IMPROVE are:

- To establish current visibility and aerosol conditions in mandatory Class I areas;
- To identify chemical species and emission sources responsible for existing man-made visibility impairment;
- To document long-term trends for assessing progress towards the national visibility goal and
- With the enactment of the regional haze rule, to provide regional haze monitoring representing all visibility-protected federal Class I areas.

Class I areas were designated based on an evaluation required by Congress in the 1977 federal Clean Air Act amendments. The evaluation, which the U.S. Forest Service and National Park Service performed, reviewed the wilderness areas of parks and national forests which were designated as wilderness before 1977, were more than 6,000 acres in size and have visual air quality as an important resource for visitors. Of the 156 Class I areas designated across the nation, 12 are located in Arizona.

The Arizona Class I visibility network consists of a combination of visibility monitoring sites established by ADEQ and those established by the IMPROVE committee. Monitoring was conducted for the following areas:

- Grand Canyon National Park – Hance Camp,
- Grand Canyon National Park – Indian Gardens,
- Petrified Forest National Park,
- Mt. Baldy Wilderness – Greer Water Treatment Plant,
- Sycamore Canyon Wilderness – Camp Raymond,
- Mazatzal/Pine Mountain Wildernesses – Ike’s Backbone,
- Sierra Ancha Wilderness – Pleasant Valley Ranger Station,
- Superstition Wilderness – Tonto National Monument,
- Superstition Wilderness – Queen Valley,
- Saguaro National Park – West Unit,
- Saguaro National Park – East Unit,
- Chiricahua National Monument – Entrance Station,
- Galiuro Wilderness – Muleshoe Ranch (Site was closed in June of 2005) ,
- Hillside (Site was closed in June of 2005),
- Organ Pipe National Monument and
- Meadview.

Each IMPROVE site includes PM<sub>2.5</sub> sampling with subsequent analysis for the fine particle mass and major aerosol species, as well as PM<sub>10</sub> sampling and mass analysis. Many of the sites also include optical monitoring with nephelometers or transmissometers and color photography to document scenic appearance.

More information about the IMPROVE procedures, sites and data can be found on the IMPROVE website at <http://vista.cira.colostate.edu/improve/>.

### ***Urban Haze Networks***

ADEQ monitors the Phoenix and Tucson metropolitan areas with a network of instruments to characterize and quantify the extent of urban haze. There are no established federal or state standards for acceptable levels of urban haze. ADEQ began studying the nature and causes of urban hazes by conducting a study in the winter of 1989-90 in Phoenix and the winter of 1992-93 in Tucson. These studies recommended long-term, year-round monitoring of visibility. In 1993, ADEQ began deploying visibility monitoring equipment in Phoenix and Tucson. These visibility monitoring data are needed to provide policymakers and the public with information, track short- and long-term trends, assess source contributions to urban haze and better evaluate the effectiveness of air pollution control strategies.

The Phoenix urban haze network includes two transmissometers (located in Phoenix and Mesa) for measuring light extinction along a fixed path length of about 3 to 5 kilometers, four nephelometers for measuring light scattering, 5 digital camera systems to record visual characteristics of the urban area, and particulate filters for quantifying and characterizing particulate matter. The Tucson urban haze network includes one transmissometer for measuring light extinction along a fixed path length of about 3-5 kilometers, 3 nephelometers for measuring light scattering, and a digital camera system operated by Pima County to record visual characteristics of the urban area. Operation of Phoenix and Tucson area urban haze particulate monitors was discontinued at the close of 2004. Data from active PM<sub>10</sub> and PM<sub>2.5</sub> samplers will be used to characterize chemical composition and seasonal variation on an as needed basis.

The website for Phoenix area visibility is <http://www.phoenixvis.net/>. The website for the Tucson camera system is <http://www.airinfonow.org/>.

### ***Photochemical Assessment Monitoring Station Monitoring***

Section 182(c)(1) of the 1990 Clean Air Act Amendments required the administrator to promulgate rules for the enhanced monitoring of ozone, oxides of nitrogen (NO<sub>x</sub>) and volatile organic compounds (VOCs) to obtain more comprehensive and representative data on ozone air pollution. Immediately following the promulgation of those rules, the affected states were to begin actions necessary to adopt and

implement a program to improve ambient monitoring activities and the monitoring of emissions of NO<sub>x</sub> and VOCs. Each state implementation plan (SIP) for the affected areas must contain commitments to implement the appropriate ambient monitoring network for such air pollutants. The subsequent revisions to 40 CFR 58 (1993) required states to establish photochemical assessment monitoring stations (PAMS) as part of their SIP monitoring networks in ozone nonattainment areas classified as serious, severe or extreme. The principal reasons for requiring the collection of additional ambient air pollutant and meteorological data are the nationwide lack of attainment of the ozone NAAQS and the need for a more comprehensive air quality database for ozone and its precursors.

The chief objective of the enhanced ozone monitoring requirements is to provide air quality data that will assist air pollution control agencies in evaluating, tracking the progress of and, if necessary, refining control strategies for attaining the ozone NAAQS. Ambient concentrations of ozone and ozone precursors are used to make attainment and nonattainment decisions, aid in tracking VOC and NO<sub>x</sub> emission reductions, better characterize the nature and extent of the ozone problem, and examine air quality trends. In addition, data from the PAMS network provide an improved database for evaluating photochemical model performance, especially for future control strategy mid-course corrections as part of the continuing air quality management process. The data are particularly useful to states in ensuring the implementation of the most cost-effective regulatory controls.

The PAMS network array for an area should be fashioned to supply measurements that will assist states in understanding and solving ozone nonattainment problems. EPA has defined its monitoring objectives with the following five site types. Type 1 Site: Upwind and Background Characterization, Type 2 and 2a Sites: Maximum Ozone Precursor Emissions Impact, Type 3 Site: Maximum Ozone Concentration, Type 4 Site: Extreme Downwind Monitoring.

PAMS data include measurements of O<sub>3</sub>, NO<sub>x</sub>, a target list of VOCs including several carbonyls, and surface and upper air meteorology. Most PAMS sites measure 56 target hydrocarbons on either an hourly or three-hour basis during the ozone season. The Type 2 sites also collect data on three carbonyl compounds (formaldehyde, acetaldehyde and acetone) during the ozone monitoring period. Included in the monitored VOC species are 10 compounds classified as hazardous air pollutants. All stations must measure O<sub>3</sub>, NO<sub>x</sub> and surface meteorological parameters on an hourly basis. ADEQ has installed four PAMS monitoring sites to date, the ADEQ Supersite (located near 17th Avenue and Campbell) in Central Phoenix (a Type 2 site); the wind profiler (upper air meteorology) site; the Queen Valley site (Type 3); and the South Phoenix site (Type 2a). A time line describing installation dates of additional sites is provided in Table 4.

*Table 4: PAMS Installation Time Line*

Type of Ozone		Proposed Installation
PAMS	Season	
Type 2	1999	Supersite - 17th Avenue and Campbell, Phoenix
Type 2a	2001	South Phoenix - Central and Broadway
Type 3	2001	Queen Valley

### **Annual Ambient Air Monitoring Network Review**

In 1999, ADEQ expanded the scope of the annual ambient air monitoring network reviews beyond the state and local air monitoring stations (SLAMS) to include all state networks. 40 CFR §58.20(d) requires states to complete and submit to EPA an annual network review.

States are required to commit to and explain the air quality surveillance systems in their state implementation plans. The air quality surveillance systems consist of various sites designated as state and local air monitoring stations (SLAMS), national air monitoring stations (NAMS) and PAMS. To provide a complete review of the air monitoring network, ADEQ chose to include additional stations classified as special purpose monitoring stations (SPM), which includes urban haze monitoring sites, IMPROVE sites, ADEQ visibility stations located in or near mandatory Class I areas, and source-oriented monitoring sites operated independently by the permittee.

The annual network review determines conformance with the requirements of 40 CFR Part 58, Appendix D (*Network Design Criteria*) and Appendix E (*Probe and Path Siting Criteria*) for sites classified as SLAMS, NAMS, PAMS and SPM. Class I monitoring sites are subject to specific siting and operational guidance developed by the IMPROVE Steering Committee. Results of the annual network review are used to determine how well the network is achieving its required air monitoring objectives, how well it meets data users' needs and how it should be modified (through termination of existing stations, relocation of stations, establishment of new stations, monitoring of additional parameters and/or changes to the sampling schedule) to continue to meet its objectives and data needs. The main purpose of the review is to improve the network so that it provides adequate, representative and useful air quality data.

In the upcoming year, ADEQ anticipates developing or refining existing network plans for the NAAQS and urban haze ambient monitoring programs that will define

specific program goals and objectives. The initial monitoring plans will use recommendations made in the annual network review and will go through a review every two to three years considering factors such as data results and completeness, site representativeness, and data representativeness. The monitoring plan review will also tabulate network review results accumulated over the prior three-year period and will recommend changes to the monitoring plans and instrument or operating requirements.

### **Monitoring Methods**

The gaseous criteria pollutants ( $\text{SO}_2$ ,  $\text{O}_3$ ,  $\text{NO}_2$  and  $\text{CO}$ ),  $\text{PM}_{10}$  at those sites with continuous instruments, and optical characteristics of the atmosphere (total light extinction, light absorption by gases, light scattering by particles and light absorption by particles), are monitored with continuous analyzers taking approximately one pollutant sample per second. These values are averaged on an hourly basis and recorded to the correct number of significant digits, based on the form of the air quality standards and the detection limits of the instrument. In most cases, the hourly data are summarized into the appropriate multi-hour averages. The agency or company network operators conduct regular checks of the stability, reproducibility, precision and accuracy of these instruments. Precision and accuracy of ambient data are assessed across an entire network using statistical tests that EPA requires.

Particulate matter,  $\text{PM}_{10}$  and  $\text{PM}_{2.5}$ , is usually sampled for 24 hours, from midnight to midnight, most often on every sixth day, with the following filter-based method. Using a timer, ambient air is drawn through an inlet of a specified design at a known flow rate onto a filter that collects all PM less than a diameter specified by the inlet design. The filters are weighed before and after the sample period to determine the difference in mass and then divided by the product of the flow rate with the elapsed time to arrive at a mass per unit volume concentration. Some filters are subjected to chemical analysis to determine the amount of various analytes and integrated with the flow rate and timer information to calculate their concentrations. These data are summarized into the appropriate quarterly or annual averages. These samplers are also certified as federal reference or equivalent methods. The agency or company network operators perform regular checks of the stability, reproducibility, precision and accuracy of the samplers and laboratory procedures. Again, precision and accuracy of ambient data are assessed across an entire network using statistical tests that EPA requires.

Visibility monitoring methods are generally divided into the three groups of optical, scene and aerosol (PM). Monitoring of visibility requires qualitative and quantitative information about the causes of haze (e.g., what is in the air, the formation, transport and deposition of pollutants) and the nature of haze (the optical effects of those pollutants to the observer). Scene conditions of visual air quality associated with hazes are recorded with a camera. In the past, ADEQ has used a super-VHS video

format and 35 mm slides. The video camera was programmed to advance at the rate of one frame every four minutes during daylight hours. When scene information is obtained from 35 mm slides, a picture is taken at the same times each day to establish baseline conditions and track variations in haze. ADEQ is currently replacing 35 mm slides with digital and Web cameras for continued documentation of scene conditions.

Quantitative measurement of light extinction ( $B_{\text{ext}}$ ) has four components:

- Light scattering by gases ( $B_{\text{sg}}$ )
- Light absorption by gases ( $B_{\text{ag}}$ )
- Light scattering by particles ( $B_{\text{sp}}$ )
- Light absorption by particles ( $B_{\text{ap}}$ )

Mathematically, the relationship is expressed as  $B_{\text{ext}} = B_{\text{sg}} + B_{\text{ag}} + B_{\text{sp}} + B_{\text{ap}}$ , where the units are inverse megameters ( $\text{Mm}^{-1}$ ), or the amount of light removed per million meters of distance a viewer looks through.

Total optical light extinction ( $B_{\text{ext}}$ ) is measured directly with a device called a transmissometer. The transmissometer generates visible light in the same wavelength (550 nanometers) as the human eye detects and then transmits that light beam over a sight path of several kilometers to a photocell detector. The transmissometer's design and operation allow its data to be directly correlated with human perception of visibility through the atmosphere. Transmissometer data are also used to check the general accuracy of the sum of the components of light extinction as measured by other continuous monitors. Transmissometers have been operated in Phoenix and Tucson since 1993.

Light scattering by gases ( $B_{\text{sg}}$ ) is a function of air density and is unrelated to air pollution sources. This parameter is derived and does not require measurement. In contrast, the other three components of light extinction are human-caused and require measurement with continuous monitors.

Light absorption by gases ( $B_{\text{ag}}$ ) is determined by continuously measuring nitrogen dioxide ( $\text{NO}_2$ ) since it is the only gas normally present in urban or Class I areas that absorbs significant quantities of visible light. Several EPA reference or equivalent method  $\text{NO}_2$  monitors are deployed to verify maintenance of the NAAQS throughout Arizona, including monitoring at Tucson, Phoenix, Queen Valley and Tonto National Monument, while the National Park Service network tracks  $\text{NO}_2$  at several national parks in Arizona.

Light scattering by particles ( $B_{\text{sp}}$ ) is determined by continuously, directly measuring particle scattering variation in a calibrated ambient sampling chamber called a nephelometer. The nephelometer samples air at ambient temperature and relative

humidity conditions. Routine monitoring with this instrument began in both the Class I area and urban haze networks during 1996. Light absorption by particles ( $B_{ap}$ ) is determined by continuously measuring the quantity of light transmitted through a filter tape or intermittently through a filter from a PM sampler. Data from these analyses are reported in micrograms per cubic meter ( $\mu\text{g}/\text{m}^3$ ) of elemental carbon and are converted to the  $B_{ap}$  units of  $\text{Mm}^{-1}$  using a laboratory-derived light absorption coefficient. Routine data collection using a continuous instrument, the aethalometer, began in December 1996 in Phoenix and February 1998 in Tucson.  $B_{ap}$  is also measured intermittently using the PM sample filters collected in both the Class I area and urban haze networks.

In monitoring visibility, it is also essential to collect and analyze particulate samples to define and to understand the chemistry of aerosols present before, during and after haze events. The chemical speciation data can be used to determine the contributions of each source category to the observed optical haze data. From these filter data, the chemical components are used to calculate light extinction for the filter sample period and compared with continuous measurements as a check. Finally, the samplers used in the urban haze networks also monitor compliance with  $\text{PM}_{10}$  and  $\text{PM}_{2.5}$  national air quality standards and provide information on the categories of pollution sources contributing to observed  $\text{PM}_{10}$  and  $\text{PM}_{2.5}$  concentrations. Sampling frequency for PM in the urban networks is generally every sixth day in the ADEQ network and every third day in the IMPROVE Class I area network. Every day sampling at all monitoring sites would be cost-prohibitive and personnel-intensive using current particulate sampling technologies.

To more fully understand the causes of hazes often associated with certain atmospheric conditions, it is necessary to monitor certain meteorological parameters. For these reasons, each network includes meteorological data such as temperature, relative humidity, wind speed and direction. Routine measurements of upper air temperature and water vapor are not made in the Phoenix area but information from the twice-daily rawinsonde launches by the National Weather Service at Tucson, Flagstaff, Las Vegas, Nevada and El Paso, Texas are used to characterize the air masses over Arizona.

## Monitoring Data

---

### Introduction

Air quality measurements in Arizona can be divided into the three categories of criteria pollutants, visibility and photochemical monitoring. Each category is discussed below.

EPA has set National Ambient Air Quality Standards (NAAQS) for the criteria air pollutants, which are CO, ozone, nitrogen dioxide, sulfur dioxide, lead and particulate matter 10 microns in size and smaller ( $PM_{10}$ ) and particulate matter 2.5 microns in size and smaller ( $PM_{2.5}$ ).

These pollutants are monitored in Arizona by industry, county air pollution districts, the National Park Service, Indian tribes and ADEQ.

The 2005 data measurements by criteria pollutant begin below. The

data tables in this section are organized by county; site operator information can be found in the site index tables in Appendix 1. Data recovery information (valid samples as a percent of total scheduled samples) is included in the tables. The number of valid samples is important for determining the representativeness of the average data calculations. Information about the compliance requirements and status for the criteria pollutants begins on Page 36. Visibility monitoring information is presented beginning on Page 58.




**Figure 3 – ADEQ's Phoenix James L. Guyton Supersite monitoring station.**

### Criteria Pollutants - 2005 Data

#### Carbon Monoxide

Carbon monoxide (CO) - a colorless, odorless, tasteless gas that is produced in the incomplete combustion of fuels - has a variety of adverse health effects that arise from its ability to chemically bind with blood hemoglobin. CO successfully competes with oxygen for binding with hemoglobin and thereby impairs oxygen transport. This impaired transport leads to several central nervous system effects, such as the impairment of time interval discrimination, changes in relative brightness thresholds, increased reaction time, and headache, fatigue and dizziness. CO exposures also contribute to or exacerbate arteriosclerotic heart disease.

In Arizona's metropolitan areas, about 51 percent of CO emissions come from on-road motor vehicles; 45 percent from off-road vehicles or equipment such as construction, lawn and garden equipment; and the remainder from point and area sources. This pollutant has low background levels, with highest concentrations next to busy streets, and has elevated neighborhood concentrations in locations that reflect emissions transported from upwind areas. Its concentrations peak from November to January because its emissions are highest in cold weather - automotive emissions of CO vary inversely with temperature - and because the surface layer of the atmosphere is at its most stable in wintertime. Hourly concentrations tend to be at their maximum during the morning rush hour and between 6 p.m. and midnight.



Controls have reduced CO emissions, and the standards have been achieved in the metropolitan Phoenix area since 1996, in stark contrast to the first half of the 1980s, when more than 100 exceedances were recorded each year. Similar improvements have occurred in Tucson, where the last eight-hour exceedances were recorded in 1988 at two sites. Equipping vehicles with catalytic converters and electronic ignition systems were the most effective controls, but significant reductions can also be attributed to the vehicle inspection program (beginning in 1976) and oxygenated fuels (beginning in 1989).

CO is monitored continuously with non-dispersive infrared instruments that are deployed in urban neighborhoods and near busy roadways or intersections. In 2005, 14 monitors were operated in greater Phoenix; 6 monitors were operated in metropolitan Tucson. Monitors in Apache Junction and Casa Grande were closed during 2002. Table 5 presents the 2005 CO data.

<b>Table 5: 2005 Carbon Monoxide Data (in ppm)</b> (NAAQS 1-hour 35 ppm, 8-hour 9 ppm)						
Site or City	One-Hour Average Value		Eight-Hour Average Value		Valid Data Recovery*	
	Max Value	2nd High	Max Value	2nd High	No. of Obs.	%
<b>Maricopa County</b>						
Buckeye <sup>S</sup>	1.1	1.1	0.9	0.9	4942	97
Central Phoenix	5.2	5.1	4.1	3.8	8556	98
Dysart <sup>S</sup>	1.7	1.7	1.3	1.2	5005	98
Glendale <sup>S</sup>	3.2	3.1	2.4	2.3	4872	96
Greenwood	5.9	5.4	4.2	4.1	8591	98
JLG Supersite	5.6	5.1	3.7	3.6	8714	99
Mesa <sup>S</sup>	3.4	3.3	2.4	2.4	4865	96
North Phoenix <sup>S</sup>	3.8	3.5	2.3	2.2	5021	99
South Phoenix <sup>S</sup>	5.5	5.2	3.8	3.2	4980	98
South Scottsdale <sup>S</sup>	3.2	3.1	2.4	2.4	4813	95
Tempe <sup>S</sup>	3.2	3.0	2.6	2.4	5000	98
West Chandler <sup>S</sup>	3.5	2.7	2.4	2.0	4998	98
West Indian School RD	6.8	6.5	5.3	4.8	8415	96
West Phoenix	7.2	7.0	5.8	4.6	8407	96
<b>Pima County</b>						
22nd St. & Alvernon	4.1	3.6	2.2	2.1	8718	99
22nd St. & Craycroft	3.5	3.3	1.7	1.5	8737	99
Cherry & Glenn <sup>S</sup>	3.8	3.4	2.5	2.4	5070	99
Children's Park	2.0	1.8	1.1	1.1	8751	99
Golf Links & Kolb <sup>S</sup>	3.3	3.2	2.2	2.1	5071	99
Tucson Downtown	3.0	2.8	1.9	1.7	8740	99

\* **Valid Data Recovery** shows the number of valid samples collected during the year and the percentage of the 8760 sampling hours in the year that were valid. Percentages will always be less than 100% due to mandatory quality assurance testing of the monitors requiring them to be off-line for several hours at a time.

<sup>S</sup> Seasonal monitor, operational during January 1 to April 1 and September 1 to December 31; 5088 sampling hours in non leap years.

Exceptions:


Pima County seasonal monitors operated January 1 - April 30 and October 1 - December 31; 5088 sampling hours in non leap years.

## *Nitrogen Dioxide*

Nitrogen dioxide (NO<sub>2</sub>) is a reddish-brown gas that is formed by the oxidation of nitric oxide (NO) -- a byproduct of all combustion. At the lowest NO<sub>2</sub> exposure levels at which adverse health effects have been detected, respiratory damage has been observed: destruction of cilia, alveolar tissue disruption and obstruction of the respiratory bronchioles. Animal studies suggest that NO<sub>2</sub> may be a causal or aggravating agent in respiratory infections. However, community exposure studies to lower ambient levels of NO<sub>2</sub> have demonstrated no significant links with respiratory symptoms or disease.

This pollutant is of greater concern in its reduction of visibility (it causes 5 percent of the visibility reduction in Phoenix) and in its contributory role in the photochemical formation of ozone.

Combustion emissions of nitrogen oxides are 95 percent nitric oxide and 5 percent NO<sub>2</sub>. Because nitric oxide is rapidly oxidized to nitrogen dioxide, nitric oxide emissions serve as a surrogate for NO<sub>2</sub>. In a recent Phoenix emissions inventory, the transportation sector dominated nitric oxide emissions: 58 percent of the emissions came from cars and trucks, 27 percent came from off-road vehicles such as trains and diesel-powered construction vehicles, and 15 percent from other sources, including power plants, biogenic emissions from soil and stationary combustion sources. Nitric oxide and NO<sub>2</sub> concentrations are highest near major roadways. Nitric oxide concentrations decrease rapidly with distance from the roadway, whereas NO<sub>2</sub> concentrations are more evenly distributed because of their formation through oxidation and their subsequent transport. Concentrations of NO<sub>2</sub> are highest in the late afternoon and early evening of winter, when rush hour emissions of nitric oxide are converted to NO<sub>2</sub> under relatively stable atmospheric conditions. Because nitric oxide reacts rapidly with ozone, nocturnal ozone concentrations in cities are often reduced to near-zero levels. This nitric oxide scavenging of ozone does not occur in remote areas. Nocturnal ozone concentrations at background sites are high compared with the urban concentrations.



Nitrogen oxides emissions from motor vehicles have been reduced through retardation of spark timing, lowering the compression ratio, exhaust gas recirculation systems and three-way catalysts. The vehicle inspection program, with its NO<sub>x</sub> test for light-duty gasoline vehicles 1981 and newer (in Phoenix only) has also helped. Reformulated gasolines also decrease nitrogen oxides emissions: Federal Phase II gasoline, by 1.5 percent for vehicular and 0.5 percent for off-road equipment; California Phase 2 gasoline, by 6.4 percent for vehicular and 7.7 percent for off road equipment.

NO<sub>2</sub> is monitored continuously with chemiluminescence instruments, which also determine nitric oxide (NO) concentrations and NO<sub>x</sub> (the sum of NO<sub>2</sub> and NO) concentrations. These instruments are located in urban neighborhoods where either the emissions are dense or where ozone concentrations tend to be at their maximum. In addition, these monitors are located near major coal-fired electrical power plants. Eleven monitors were operated in Arizona in 2005. Table 6 presents the NO<sub>2</sub> data available in 2005.

<b>Table 6: 2005 Nitrogen Dioxide (in ppm)</b> (NAAQS Annual Mean 0.053 ppm)				
Site or City	Annual Average	Maximum Value	Valid Data Recovery *	
		One-Hour Average	No. of Obs.	%
<b>Apache County</b>				
Springerville – Coyote Hills	.0013	.014	8573	98
<b>La Paz</b>				
Alamo Lake <sup>S</sup> (Opened 05/20/05)	.0024	.011	3091	96
<b>Maricopa County</b>				
Buckeye	.0119	.053	8307	95
Central Phoenix	.0262	.095	8490	97
Greenwood	.0315	.131	8467	97
JLG Supersite <sup>S</sup>	.0208	.077	5096	99
South Scottsdale	.0196	.079	8424	96
West Phoenix	.0235	.100	8191	94
<b>Pima County</b>				
22nd St. & Craycroft	.0149	.056	8681	99
Children's Park	.0152	.049	8655	99
<b>Yuma County</b>				
Yuma Game & Fish <sup>S</sup>	.0103	.051	6549	99

\* **Valid Data Recovery** shows the number of valid observations and the percentage of the possible 8760 hourly samples during the year (always be less than 100% due to mandatory quality assurance testing requiring the monitors to be off-line for several hours at a time).

<sup>S</sup> Seasonal Monitors:


Phoenix JLG Supersite operates during winter CO season, January 1 to April 30 and October 1 to December 31; 5112 hours in 2005.

Yuma Game & Fish operated April 1 to the end of 2005; 6600 hours in 2005.

## Sulfur Dioxide

Exposure to sulfur dioxide (SO<sub>2</sub>), a colorless gas with a pungent, irritating odor at elevated concentrations, alters the mechanical function of the upper airway, including increasing the nasal flow resistance and decreasing the nasal mucus flow rate. Short-term exposures result in an exaggerated air flow resistance in about 10 percent of the subjects tested and produce acute bronchioconstriction in strenuously exercising asthmatics.

In Arizona, the principal source of SO<sub>2</sub> emissions has been the smelting of sulfide copper ore. Most fuels contain trace quantities of sulfur, and their combustion releases both gaseous SO<sub>2</sub> and particulate sulfate (SO<sub>4</sub><sup>2-</sup>). A recent emissions inventory for Phoenix shows 32 percent of SO<sub>2</sub> emissions come from point sources, 26 percent from area sources, 23 percent from off-road vehicles and equipment, and 19 percent from on-road motor vehicles. SO<sub>2</sub> is removed from the atmosphere through dry deposition on plants and its conversion to sulfuric acid and eventually to sulfate. SO<sub>2</sub> has extremely low background levels, with elevated concentrations found downwind of large point sources. Concentrations in urban areas are low and are homogeneously distributed, with annual averages varying from 3 to 10 µg/m<sup>3</sup>, well within the annual standard of 80 µg/m<sup>3</sup>.



Major controls were installed in Arizona's copper smelters in the 1980s, which reduced SO<sub>2</sub> emissions substantially. Vehicular emissions of SO<sub>2</sub> and sulfate have been reduced through lowering the sulfur content in diesel fuel and gasoline.

SO<sub>2</sub> is monitored continuously with pulsed fluorescence instruments, most of which are clustered around copper smelters or coal-fired electric power plants. In 2005, ten reporting monitors were sited near copper smelters, one near a power plant and four in urban areas. Table 7 presents the SO<sub>2</sub> data collected in Arizona in 2005 from the monitors near copper smelters and in urban areas.


**Table 7: 2005 Sulfur Dioxide (in  $\mu\text{g}/\text{m}^3$ )**  
 (Primary NAAQS Annual Average  $80 \mu\text{g}/\text{m}^3$ [0.030 ppm], 24-hour Average  $365 \mu\text{g}/\text{m}^3$ [0.14 ppm]  
 Secondary NAAQS 3-hour  $1300 \mu\text{g}/\text{m}^3$  [0.5 ppm])

Site or City	Annual Average	Maximum Value				Valid Data Recovery *	
		3-Hour Average		24-Hour Average		No. Obs.	%
		Max Value	2nd High	Max Value	2nd High		
<b>Apache County</b>							
TEP – Springerville – Coyote Hills	1	21	19	16	11	8480	97
<b>Gila County</b>							
ASARCO – Globe Hwy.	34	873	753	182	152	8616	98
ASARCO – Hayden – Garfield AVE	18	691	621	197	152	8667	99
ASARCO – Montgomery Ranch	35	596	575	210	175	8635	99
Hayden– Old Jail, ADEQ	22	806	603	157	113	8703	99
Hayden– Old Jail, ASARCO	16	744	616	139	82	8649	99
Miami – Ridgeline	12	252	238	79	76	8686	99
PDMI – Miami – Jones Ranch	14	395	307	95	87	8738	99
PDMI –Miami–Town Site	10	273	221	59	54	8752	99
<b>Maricopa County</b>							
Central Phoenix	6	55	31	21	18	8589	98
JLG Supersite # (Opened 03/01/2005)	7	21	21	16	16	7235	99
South Scottsdale	4	18	16	16	16	8380	96
<b>Pima County</b>							
22nd St. & Craycroft	3	26	24	10	8	8682	99
<b>Pinal County</b>							
ASARCO - Hayden Jct.	10	250	239	44	41	8655	99
San Manuel	5	16	16	8	8	8716	99

\* **Valid Data Recovery** shows the number of valid samples collected during the year and the percentage of the 8760 sampling hours in the year that were valid. Percentages will always be less than 100% due to mandatory quality assurance testing of the monitors requiring them to be off-line for several hours at a time. Exceptions: JLG Supersite operated March 1 - December, 7344 sampling hours. Note: Sulfur dioxide conversion factor: ppm x 2620 =  $\mu\text{g}/\text{m}^3$ . # Less than 75% data recovery; does not satisfy EPA criteria.

## Ozone

Ozone ( $O_3$ ) - a colorless, slightly odorous gas - is both a natural component of the atmosphere, through its photochemical formation from natural sources of CO, hydrocarbons and nitrogen oxides, and an important air contaminant in urban atmospheres. In the stratosphere,  $O_3$  blocks harmful ultraviolet radiation. In the urban atmosphere, its formation from anthropogenic emissions of hydrocarbons and nitrogen oxides leads to concentrations harmful to people, animals, plants and materials.  $O_3$  causes significant physiological and pathological changes in both animals and humans at concentrations present in many urban environments. Short-term (one to two hours) exposures to concentrations in the range of 0.1 to 0.4 parts per million induce changes in lung function, including increased respiratory rates, increased pulmonary resistance, decreased tidal volumes and changes in lung mechanics. Symptomatic responses in exercising adults include throat dryness, chest tightness, substernal pain, cough, wheeze, pain on deep inspiration, shortness of breath and headache. These symptoms also have been observed at lower concentrations for longer exposures. Evidence suggests that  $O_3$  exposure makes the respiratory airways more susceptible to other bronchioconstrictive challenges. Animal studies suggest that ozone exposure interferes with or inhibits the immune system.  $O_3$  at ambient concentrations injures the stomates, which are the cells that regulate plant respiration, resulting in flecks on the upper leaf surfaces of dichotomous plants and the death of the tips of coniferous needles.  $O_3$  is considered by plant scientists to be the most important of all of the phytotoxic air pollutants, causing over 90 percent of all plant injury from air pollution on a global basis.



$O_3$ , formed photochemically by the reaction of volatile organic compounds and nitrogen oxides, has elevated concentrations only in the summer. Volatile organic compound (VOC) emissions in greater Phoenix come from cars and trucks (31 percent), off-road vehicles and equipment such as lawn mowers (27 percent), small stationary sources (20 percent), biogenic emissions from grass, shrubs and trees (17 percent) and point sources (5 percent). NO<sub>x</sub> comes from cars and trucks (58 percent), off-road vehicles such as construction equipment and trains (27 percent), electric power plants (7 percent), small stationary sources (4 percent) and biogenic emissions from soil (4 percent).  $O_3$  has relatively high background levels, with the daily maximum in remote areas being about one-half to three-quarters of the daily maximum in the urban areas. In an urban area, the highest  $O_3$  concentrations tend to occur on the downwind edge, although high concentrations do occur less frequently in the central city. High  $O_3$  concentrations are a summer phenomenon caused when sunlight, biogenic emissions, and evaporative hydrocarbon emissions peak. Urban  $O_3$  concentrations are low to near zero at night, rise rapidly through the morning and peak in the afternoon.

Controls to reduce the precursors of ozone - VOC and NO<sub>x</sub> - have been successfully implemented for years. NO<sub>x</sub> and VOC from vehicular exhaust have been reduced through engine modifications and three-way catalytic converters. Evaporative hydrocarbons from vehicles have been reduced through better engineered fuel tanks and auxiliary plumbing combined with carbon absorption canisters. Additional reductions of vehicular VOC have come through ADEQ's vehicle inspection program, which tests all gasoline vehicles for hydrocarbons (Phoenix and Tucson), through vapor-capturing equipment for gasoline tankers, vapor recovery systems at retail gas stations (Phoenix area only) and cleaner burning gasoline (Phoenix area only). Stationary source hydrocarbons have been reduced through a variety of better control equipment required by stricter regulations. Despite these efforts, the continued population growth in Arizona combined with the high natural background O<sub>3</sub>, may make achieving the eight-hour standard difficult.

Ultraviolet absorption instruments monitor O<sub>3</sub> continuously in urban neighborhoods for population exposure, in areas downwind of urban areas for maximum concentration monitoring and in remote areas for background information. In 2005, 35 reporting O<sub>3</sub> monitors were in operation; five for background, 25 for urban neighborhoods and 10 for maximum concentrations downwind of urban areas. Tables 8 and 9 present the 2005 Arizona O<sub>3</sub>.

<b>Table 8: 2005 Ozone Data (in ppm), One-Hour Averages</b> (NAAQS 1-hour 0.12 ppm)						
Site or City	Max Value	2nd High	3rd High	4th High	Valid Data Recovery*	
					No. Of Days	%
<b>Cochise County</b>						
Chiricahua NM Entrance	.077	.076	.076	.074	348	95
<b>Coconino County</b>						
Grand Canyon NP Hance	.093	.089	.086	.083	360	99
<b>Gila County</b>						
Tonto NM <sup>S</sup>	.113	.109	.108	.101	213	99
<b>La Paz</b>						
Alamo Lake <sup>S</sup> # (Opened 05/20/05)	.081	.080	.077	.075	130	61
<b>Maricopa County</b>						
Blue Point	.107	.105	.098	.093	365	100
Buckeye <sup>S</sup>	.085	.080	.076	.075	214	100

**Table 8: 2005 Ozone Data (in ppm), One-Hour Averages  
(NAAQS 1-hour 0.12 ppm)**

Site or City	Max Value	2nd High	3rd High	4th High	Valid Data Recovery*	
					No. Of Days	%
Cave Creek <sup>S</sup>	.108	.093	.092	.092	213	99
Central Phoenix	.099	.092	.092	.090	360	99
Dysart <sup>S</sup>	.082	.081	.080	.078	214	100
Falcon Field <sup>S</sup>	.104	.095	.093	.091	206	96
Fountain Hills	<b>.129</b>	.115	.106	.106	360	99
Glendale <sup>S</sup>	.096	.093	.090	.088	214	100
Humboldt Mt. <sup>S</sup>	.104	.099	.096	.094	204	95
JLG Supersite	.095	.094	.093	.093	365	100
North Phoenix	.110	.109	.108	.102	363	99
Pinnacle Peak	.104	.100	.094	.094	359	98
Rio Verde <sup>S</sup>	.117	.114	.110	.109	213	99
South Phoenix	.108	.096	.094	.092	362	99
South Scottsdale	.117	.100	.099	.094	359	98
Tempe <sup>S</sup>	.111	.104	.099	.097	214	100
West Chandler <sup>S</sup>	.096	.091	.088	.086	214	100
West Phoenix	.094	.086	.081	.081	344	96
<b>Navajo County</b>						
Petrified Forest NP	.101	.094	.080	.076	335	92
<b>Pima County</b>						
22nd & Craycroft	.093	.089	.089	.084	364	99
Children's Park	.086	.085	.084	.083	365	100
Coachline	.075	.073	.073	.073	365	100
Green Valley	.081	.079	.079	.074	361	99
Rose Elementary	.075	.074	.074	.073	365	100
Saguaro National Park East	.101	.091	.090	.089	365	100
Tangerine	.080	.079	.079	.077	365	100
Tucson Downtown	.084	.083	.079	.078	365	100
Tucson Fairgrounds	.085	.085	.083	.083	353	97
<b>Pinal County</b>						
Apache Junction Maintenance Yard	.097	.089	.085	.083	358	98
Casa Grande Airport	.089	.088	.081	.080	358	98

**Table 8: 2005 Ozone Data (in ppm), One-Hour Averages  
(NAAQS 1-hour 0.12 ppm)**

Site or City	Max Value	2nd High	3rd High	4th High	Valid Data Recovery*	
					No. Of Days	%
Queen Creek <sup>S##</sup>	.093	.091	.088	.087	213	99
Maricopa <sup>S</sup>	.079	.078	.075	.069	211	99
Pinal Air Park <sup>S</sup>	.088	.085	.084	.084	207	97
Queen Valley <sup>S</sup>	.117	.113	.110	.105	214	100
<b>Yavapai County</b>						
Hillside <sup>S #</sup> (Closed 06/04/2005)	.078	.078	.078	.078	58	27
<b>Yuma County</b>						
Yuma Game & Fish <sup>S</sup>	.090	.089	.085	.085	214	100

\* **Valid Data Recovery** shows the number of days with at least 75 percent (18 or more hours) of valid data recovery. It also shows the percentage of the total number of scheduled sampling days that meet that criterion. Scheduled sampling days for non-seasonal monitors in 2005 was 365.

<sup>S</sup> Seasonal monitor, operational during April 1 to November 1; 214 scheduled sampling days in the season.

# Less than 75% data recovery; does not satisfy EPA summary criteria.

## Formerly "Combs"

**Table 9: 2005 Ozone Data (in ppm), Eight-Hour Averages  
(NAAQS 8-hour 0.08 ppm)**

Site or City	Max Value	2nd High	3rd High	4th High	Daily Exceed.	Valid Data Recovery *	
						No. of Days	%
<b>Cochise County</b>							
Chiricahua NM Entrance	.073	.073	.072	.072	0	346	95
<b>Coconino County</b>							
Grand Canyon NP Hance	.089	.083	.080	.079	1	355	97
<b>Gila County</b>							
Tonto NM <sup>S</sup>	.098	.097	.084	.084	2	213	99
<b>La Paz</b>							
Alamo Lake <sup>S</sup> # (Opened 05/20/05)	.076	.075	.072	.071	0	127	59
<b>Maricopa County</b>							
Blue Point	.089	.088	.083	.081	2	364	99
Buckeye <sup>S</sup>	.067	.066	.066	.065	0	214	100
Cave Creek <sup>S</sup>	.084	.083	.083	.082	0	213	99
Central Phoenix	.081	.080	.078	.075	0	357	98
Dysart <sup>S</sup>	.073	.069	.067	.066	0	213	99
Falcon Field <sup>S</sup>	.081	.078	.078	.076	0	201	94
Fountain Hills	.096	.091	.088	.088	6	360	99
Glendale <sup>S</sup>	.078	.077	.076	.076	0	213	99
Humboldt Mt. <sup>S</sup>	.088	.088	.087	.087	5	201	94
JLG Supersite	.079	.077	.076	.076	0	365	100
North Phoenix	.089	.088	.085	.084	3	362	99
Pinnacle Peak	.085	.083	.083	.083	1	357	99
Rio Verde <sup>S</sup>	.093	.088	.087	.087	6	212	99
South Phoenix	.081	.081	.076	.076	0	359	98
South Scottsdale	.089	.084	.079	.077	1	353	97
Tempe <sup>S</sup>	.086	.078	.077	.076	1	214	100
West Chandler <sup>S</sup>	.082	.076	.075	.075	0	213	99
West Phoenix	.072	.071	.069	.068	0	338	93
<b>Navajo County</b>							
Petrified Forest NP	.082	.081	.072	.070	0	308	84

**Table 9: 2005 Ozone Data (in ppm), Eight-Hour Averages**  
(NAAQS 8-hour 0.08 ppm)

Site or City	Max Value	2nd High	3rd High	4th High	Daily Exceed.	Valid Data Recovery *	
						No. of Days	%
<b>Pima County</b>							
22nd & Craycroft	.083	.079	.075	.074	0	364	99
Children's Park	.079	.076	.075	.075	0	365	100
Coachline	.070	.067	.067	.066	0	364	99
Green Valley	.075	.074	.073	.068	0	358	98
Rose Elementary	.069	.068	.068	.067	0	365	100
Saguaro NP East	.087	.082	.080	.079	1	365	100
Tangerine	.077	.073	.073	.073	0	365	100
Tucson Downtown	.077	.071	.070	.070	0	365	100
Tucson Fairgrounds	.077	.074	.073	.073	0	350	96
<b>Pinal County</b>							
Apache Junction Maintenance Yard	.076	.073	.070	.068	0	356	97
Casa Grande Airport	.081	.074	.073	.072	0	356	97
Queen Creek <sup>S##</sup>	.080	.076	.069	.067	0	212	99
Maricopa <sup>S</sup>	.069	.069	.067	.061	0	208	97
Pinal Air Park <sup>S</sup>	.079	.079	.077	.077	0	207	97
Queen Valley <sup>S</sup>	.096	.095	.085	.084	3	214	100
<b>Yavapai County</b>							
Hillside <sup>S #</sup> (Closed 6/04/2005)	.076	.076	.076	.074	0	58	27
<b>Yuma County</b>							
Yuma Game & Fish <sup>S</sup>	.082	.079	.078	.078	0	214	100

\* **Valid Data Recovery** shows the number of days with at least 75 percent (18 or more hours) of valid data recovery. It also shows the percentage of the total number of scheduled sampling days that meet that criterion.

Scheduled sampling days for non-seasonal monitors in 2005 was 365.


<sup>S</sup> Seasonal monitor, operational during April 1 to November 1; 214 scheduled sampling days in the season.

# Less than 75% data recovery; does not satisfy EPA summary criteria.


## Formerly "Combs"

### *Particulate Matter Smaller Than 10 Microns (PM<sub>10</sub>) and Smaller Than 2.5 Microns (PM<sub>2.5</sub>)*

Particulate matter is a collective term describing very small solid or liquid particles that vary considerably in size, geometry, chemical composition and physical properties. Produced by both natural processes (pollen and wind erosion) and human activity (soot, fly ash, and dust from paved and unpaved roads), particulates contribute to visibility reduction, pose a threat to public health and cause economic damage through soil disturbance. Some fine particulates (PM<sub>2.5</sub>) are formed by the condensation of vapors or by their subsequent growth through coagulation or agglomeration. Others are emitted directly from the sources, either by combustion or from mechanical grinding of soils. Coarse particulates (2.5 to 10 microns) are formed through mechanical processes such as the grinding of matter and the atomization of liquids. Fine particulates can also be classified as primary - produced within and emitted from a source with little subsequent change - or secondary - formed in the atmosphere from gaseous emissions. Secondary particulate nitrates and sulfates, for example, form in the atmosphere from the oxidation of gaseous SO<sub>2</sub> and NO<sub>2</sub>. In contrast, most atmospheric carbon is primary, having been emitted directly from combustion sources, although some of the organic carbon in the aerosol is secondary, having been formed by the complex photochemistry of gaseous volatile organic compounds.



The size, shape and chemical composition of particulates determine their health effects. Particles larger than 10 microns are deposited in the upper respiratory tract. Particles from 2.5 to 10 microns are inhalable and are deposited in the upper parts of the respiratory system. Particles smaller than 2.5 microns are respirable and enter the pulmonary tissues to be deposited there. Particles in the size range of 0.1 to 2.5 microns are most efficiently deposited in the alveoli, where their effective toxicity is greater than larger particles because of the higher relative content of toxic heavy metals, sulfates and nitrates. Epidemiological studies have shown causal relationships between particulates and excess mortality, aggravation of bronchitis, and, in children, small, reversible changes in pulmonary function. Acidic aerosols have been linked to the inability of the upper respiratory tract and pulmonary system to remove harmful particles.



The Arizona Comparative Environmental Risk Project - a multi-disciplinary investigation into human exposure to all environmental risks completed in 1995 - ranked outdoor air quality in general and particulate matter in particular as the highest environmental risk in the state. In this study, annual premature deaths from exposure to PM<sub>10</sub> concentrations in Arizona were estimated at 963, which included 667 in Maricopa County and 88 in Tucson.

Increased percentages of hospital admissions for respiratory disease (1 to 4 percent, depending on the city), of asthma episodes (5 to 14 percent), of lower respiratory symptoms (5 to 15 percent) and of coughs (2 to 6 percent) were attributed to the prevailing annual PM<sub>10</sub> concentrations in 1991. Chronically high particulate concentrations in the ambient air continue to pose a serious health threat to many Arizonans.

Coarse particulate emissions are mostly geological and are dominated by dusts from three activities: re-entraining dust from paved roads, driving on unpaved roads and earthmoving associated with construction. Soil dust from these sources and others contribute more than 70 percent of the coarse particulates in Phoenix. On days with winds in excess of 15 miles per hour, wind erosion of soil contributes to this loading. With a more diverse chemical composition, fine particulate (PM<sub>2.5</sub>) emissions are more evenly distributed among a larger number of sources. At the Phoenix JLG Supersite, receptor modeling indicates gasoline and diesel engine exhaust account for more than two-thirds of the PM<sub>2.5</sub> emissions. Soil dust contributes another 10.5 percent.

In other urban and rural areas, this mixture of sources will vary. Agricultural and mining areas, for example, will be more heavily influenced by emissions from these activities.

PM<sub>2.5</sub> concentrations tend to be at their highest in the central portions of urban areas, diminishing to background levels at the urban fringe. In contrast, PM<sub>10</sub> concentrations are not spatially distributed smoothly because each monitoring site is strongly influenced by the degree of localized emissions of coarse particulates. Background concentrations of PM<sub>10</sub> are about 40 percent of the urban maxima (20 µg/m<sup>3</sup> for an annual average background versus about 50 µg/m<sup>3</sup> for the urban maximum). Background concentrations of PM<sub>2.5</sub> are about 5 µg/m<sup>3</sup>, in contrast to the urban maxima of 12 to 15 µg/m<sup>3</sup>. Concentrations of both size ranges of particulates tend to be higher in the late fall and winter, when atmospheric dispersion is at a seasonal low. PM<sub>10</sub> maximum concentrations can occur in any season, provided nearby sources of coarse particulates are present or when strong and gusty winds suspend soil disturbed by human activities. Hourly concentrations of particulates tend to peak during those hours of the worst dispersion, which is from sunset to mid-morning.

Controls to reduce particulates have been in place for decades, beginning with an ordinance that required watering to reduce dust from construction in Pima County in the 1960s. Maricopa County's umbrella dust abatement rule, Rule 310, has been revised many times through the years and now regulates construction dust, track-out dust from construction sites, and dust from unpaved parking and vacant lots. Efforts to reduce dust resuspended from paved roads have concentrated on eliminating track-out from construction sites, curbing and stabilizing road shoulders, and investigating more efficient street sweepers. Secondary fine particulates have been reduced by vehicular emission controls, which have reduced their precursor gases. Reducing gaseous hydrocarbon emissions, for example, has led to reductions in ambient concentrations of secondary organic carbon. In Maricopa County, the

Governor's Agricultural Best Management Practices Committee developed a rule containing best management practices for agricultural activities to reduce particulate emissions from tilling and harvesting activities of cropland and non-cropland. In a recent PM<sub>10</sub> SIP, the Maricopa Association of Governments committed to implement 77 new measures, including enhanced enforcement of the county dust rules, implementation of agricultural best management practices, diesel engine replacement and retirement programs and requirements for cleaner burning fireplaces.

Particulates are monitored by pulling ambient air through a filter, generally for 24 hours every sixth day, weighing the filter before and after, and measuring the volume of air sampled. The monitoring instruments are fitted with different aerodynamic devices to segregate particle size fractions. Particulates can also be monitored continuously with a tapered element oscillating microbalance (TEOM) instrument or a beta attenuation mass monitor (BAM) which utilizes a beam sensing through a paper tape.

The 2005 PM<sub>10</sub> data reported in Table 10 represent 59 monitors throughout Arizona and two in Mexico, located in Agua Prieta and Nogales, Sonora. TEOM data are included for those sites in the Phoenix metropolitan area that were required to change to everyday monitoring from every sixth day. BAM data are included for sites in Pima County. Data from collocated monitors are included if available. The data are reported in standard conditions adjusted to 25°C and 1 atmosphere pressure) as required by EPA.

EPA began a nationwide program to measure PM<sub>2.5</sub> using federal reference method monitors in anticipation of a new federal standard for fine particulates in 1999. Eleven federal reference method samplers were located in Arizona. The fine particulate portion of the PM<sub>10</sub> measurement made by dichot monitors has been measured for many years in Arizona and has served as an approximation for the PM<sub>2.5</sub> measurement; however it is not exactly equivalent to that measurement. Table 11 lists only the federal reference method measurements for 2005. The data are reported in ambient conditions (local temperature and pressure) as required by EPA. Particulate data from the IMPROVE network are not included. In 2006, the EPA changed the PM<sub>2.5</sub> NAAQS for 24hours from 65 ug/m<sup>3</sup> to 35 ug/ m<sup>3</sup>, with the effective date of December 16, 2006. The EPA also eliminated the annual standard for PM<sub>10</sub> but retained the 24-hour standard of 150 ug/m<sup>3</sup>.

<b>Table 10: 2005 PM<sub>10</sub> Data (in µg/m<sup>3</sup>)</b>						
<b>(NAAQS Annual Average 50 µg/m<sup>3</sup>, 24-hour Average 150 µg/m<sup>3</sup>)</b>						
<b>Bold denotes an exceedance, defined as any daily value greater than 150 µg/m<sup>3</sup> after rounding to the nearest 10 µg/m<sup>3</sup> and any annual average value greater than 50 µg/m<sup>3</sup> when rounded to the nearest 1 µg/m<sup>3</sup>.</b>						
Site or City	Method	Annual Average	24-Hour Average		Valid Data Recovery *	
			Max Value	2nd High	No. of Obs.	%
<b>Apache County</b>						
TEP – Springerville – Coalyard	TEOM	15	<b>213</b>	<b>198</b>	355	99
TEP – Springerville – Coyote Hills	TEOM	10	33	29	356	99
<b>Cochise County</b>						
Douglas Red Cross	Partisol	35	86	82	58	95
Paul Spur Chemical Lime Plant (1)#	Partisol	28	76	68	53	87
Paul Spur Chemical Lime Plant (2)	Partisol	28	91	72	56	92
<b>Coconino County</b>						
Flagstaff Middle School #	Partisol	17	38	35	49	80
Sedona Post Office #	Partisol	12	34	25	54	89
<b>Gila County</b>						
Hayden – Old Jail, ADEQ #	Partisol	30	124	63	56	92
PDMI – Miami – Golf Course	Dichot	21	40	39	56	92
Miami – Ridgeline, PDMI	Dichot	12	23	23	60	98
Payson Well Site #	Partisol	22	81	47	49	80
<b>Graham County</b>						
Safford #	Dichot	21	50	46	53	87
<b>Maricopa County</b>						
Bethune Elementary School (1)	Dichot	59	<b>198</b>	136	57	93
Bethune Elementary School (2) (Closed 06/27/2005) #	Dichot	44	91	86	25	41
Buckeye	TEOM	53	<b>169</b>	<b>158</b>	360	99
Central Phoenix	TEOM	37	116	104	361	99
Central Phoenix	Hi-Vol	39	125	76	58	95

<b>Table 10: 2005 PM<sub>10</sub> Data (in µg/m<sup>3</sup>)</b>						
<i>(NAAQS Annual Average 50 µg/m<sup>3</sup>, 24-hour Average 150 µg/m<sup>3</sup>)</i>						
<b>Bold</b> denotes an exceedance, defined as any daily value greater than 150 µg/m <sup>3</sup> after rounding to the nearest 10 µg/m <sup>3</sup> and any annual average value greater than 50 µg/m <sup>3</sup> when rounded to the nearest 1 µg/m <sup>3</sup> .						
Site or City	Method	Annual Average	24-Hour Average		Valid Data Recovery *	
			Max Value	2nd High	No. of Obs.	%
Chandler	Hi-Vol	49	130	115	60	98
Durango Complex	TEOM	<b>66</b>	<b>206</b>	<b>200</b>	361	99
Dysart	Hi-Vol	29	76	68	61	100
Glendale	Hi-Vol	29	84	56	61	100
Greenwood	Hi-Vol	52	<b>173</b>	95	60	98
Higley	TEOM	51	142	121	360	99
JLG Supersite (Opened 01/01/2005)	Partisol	32	138	100	58	95
Mesa	Hi-Vol	30	86	55	60	98
North Phoenix	Hi-Vol	30	81	72	61	100
South Phoenix	Hi-Vol	55	147	107	61	100
South Scottsdale	Hi-Vol	34	121	96	61	100
West Chandler	Hi-Vol	34	94	68	60	98
West Forty Third	TEOM	<b>74</b>	<b>233</b>	<b>200</b>	362	99
West Phoenix	Hi-Vol	45	<b>155</b>	103	60	98
<b>Mohave County</b>						
Bullhead City #	Partisol	19	48	48	56	92
<b>Navajo County</b>						
Show Low #	Partisol	14	37	25	54	89
<b>Pima County</b>						
Ajo	Partisol	23	45	43	53	87
Broadway & Swan	Hi-Vol	24	46	44	60	98
Corona De Tucson	Hi-Vol	15	33	31	59	97
Green Valley <sup>2</sup>	BAM	17	54	51	358	98

<b>Table 10: 2005 PM<sub>10</sub> Data (in µg/m<sup>3</sup>)</b>						
<i>(NAAQS Annual Average 50 µg/m<sup>3</sup>, 24-hour Average 150 µg/m<sup>3</sup>)</i>						
<b>Bold</b> denotes an exceedance, defined as any daily value greater than 150 µg/m <sup>3</sup> after rounding to the nearest 10 µg/m <sup>3</sup> and any annual average value greater than 50 µg/m <sup>3</sup> when rounded to the nearest 1 µg/m <sup>3</sup> .						
Site or City	Method	Annual Average	24-Hour Average		Valid Data Recovery *	
			Max Value	2nd High	No. of Obs.	%
Orange Grove <sup>1</sup>	Hi-Vol	29	98	74	364	100
Prince Road #	Hi-Vol	37	88	88	52	85
Rillito, ADEQ	Partisol	39	84	78	59	97
Rillito, APCC	Hi-Vol	27	57	50	59	98
Santa Clara	Hi-Vol	27	82	55	59	97
South Tucson <sup>1</sup>	Hi-Vol	30	73	73	363	99
Tangerine	Hi-Vol	19	37	35	57	93
<b>Pinal County</b>						
Apache Junction Fire Station	Hi-Vol	20	47	38	59	97
Casa Grande Downtown	Hi-Vol	31	79	72	58	95
Coolidge Maintenance Yard	Hi-Vol	36	81	73	56	92
Eloy City Complex	Hi-Vol	33	73	72	59	97
Mammoth County Complex	Hi-Vol	14	33	28	61	100
Pinal Air Park	Hi-Vol	22	122	44	61	100
Pinal County Housing Complex (1)#	Hi-Vol	<b>57</b>	<b>158</b>	114	44	72
Pinal County Housing Complex (2)	Hi-Vol	<b>59</b>	<b>179</b>	<b>172</b>	61	100
Riverside Maintenance Yard	Hi-Vol	18	35	34	58	95
Stanfield	Hi-Vol	52	<b>173</b>	143	59	97
<b>Santa Cruz County</b>						
Nogales Post Office	Partisol	<b>57</b>	<b>280</b>	<b>205</b>	59	97
<b>Yavapai County</b>						
Clarkdale – NW (2)	Dichot	15	32	31	58	95
Clarkdale – SE (1)	Dichot	22	43	42	58	95

**Table 10: 2005 PM<sub>10</sub> Data (in µg/m<sup>3</sup>)**  
(NAAQS Annual Average 50 µg/m<sup>3</sup>, 24-hour Average 150 µg/m<sup>3</sup>)  
**Bold** denotes an exceedance, defined as any daily value greater than 150 µg/m<sup>3</sup> after rounding to the nearest 10 µg/m<sup>3</sup> and any annual average value greater than 50 µg/m<sup>3</sup> when rounded to the nearest 1 µg/m<sup>3</sup>.

Site or City	Method	Annual Average	24-Hour Average		Valid Data Recovery *	
			Max Value	2nd High	No. of Obs.	%
Prescott Valley #	Partisol	15	53	31	52	85
<b>Yuma County</b>						
Yuma Courthouse (1)	Partisol	35	94	86	57	93
Yuma Courthouse (2) #	Partisol	33	99	80	56	92
<b>Mexico</b>						
Agua Prieta Fire Station	Dichot	<b>68</b>	<b>172</b>	154	60	98
Sonora Nogales Fire Station	Dichot	63	<b>240</b>	<b>194</b>	60	98

<sup>1</sup> Samples collected every day - 365 sample days in 2005.

<sup>2</sup> Samples collected every hour - 8760 sample hours in 2005.

<sup>3</sup> Samples changed from every 6th day with a Hi-Vol sampler to every hour with a TEOM.

(1) Indicates the Primary monitor (used for NAAQS compliance) in a collocated pair of monitors.

(2) Indicates the Secondary monitor (used for precision and accuracy) in a collocated pair of monitors.

\***Valid data recovery** shows the number of valid observations during 2005 and the percentage of scheduled samples that were valid. There were 61 monitoring days scheduled in 2005 for monitors on the every 6th day schedule. Rillito - APCC was the only site following the every 3rd day schedule (122 observations in 2005).

# Indicates the data do not satisfy EPA's summary criteria, usually meaning less than 75 percent valid data recovery available in one or more calendar quarters.

Exceedances due to Natural Events are excluded from annual statistics.

<b>Table 11: 2005 PM<sub>2.5</sub> Data (in <math>\mu\text{g}/\text{m}^3</math>)</b>						
<b>(NAAQS Annual Average 15 <math>\mu\text{g}/\text{m}^3</math>, 24-hour Average 65 <math>\mu\text{g}/\text{m}^3</math>)</b>						
City or Site	Method	Annual Average	24-Hour Avg		Valid Data Recovery *	
			Max	2nd High	No. of Obs.	%
<b>Cochise County</b>						
Douglas Red Cross <sup>2</sup>	FRM	7.3	17.2	16.0	60	99
<b>Coconino County</b>						
Flagstaff Middle School <sup>2</sup>	FRM	6.0	18.9	12.7	55	90
<b>Gila County</b>						
Payson Well Site <sup>2</sup> #	FRM	8.3	32.2	22.9	55	90
<b>Maricopa County</b>						
JLG Supersite <sup>3</sup>	FRM	9.7	32.7	30.3	121	100
Mesa (Opened 04/28/05)	FRM	8.9	17.8	17.5	81	98
South Phoenix (Opened 01/01/2005)	FRM	12.8	56.7	40.3	118	97
West Phoenix <sup>3</sup> (1)	FRM	12.9	49.2	44.4	119	98
West Phoenix <sup>3</sup> (2)	FRM	11.8	49.9	48.3	118	97
<b>Pima County</b>						
Children's Park <sup>3</sup>	FRM	5.9	13.2	11.4	115	94
Coachline <sup>4</sup>	BAM <sup>++</sup>	7.5	22.2	21.4	8672	99
Geronimo <sup>4</sup>	BAM <sup>++</sup>	8.9	28.3	26.6	8585	98
Green Valley <sup>4</sup>	BAM <sup>++</sup>	3.7	16.4	13.7	8585	98
Orange Grove <sup>1</sup>	FRM	6.3	16.1	13.7	326	89
Rose Elementary <sup>4</sup>	BAM <sup>++</sup>	6.7	19.0	17.5	8585	99
<b>Pinal County</b>						
Apache Junction Fire Station <sup>3</sup>	FRM	5.5	12.7	10.9	111	91
Casa Grande Downtown <sup>2</sup>	FRM	7.3	19.3	16.9	53	87

<b>Table 11: 2005 PM<sub>2.5</sub> Data (in <math>\mu\text{g}/\text{m}^3</math>)</b> (NAAQS Annual Average 15 $\mu\text{g}/\text{m}^3$ , 24-hour Average 65 $\mu\text{g}/\text{m}^3$ )						
City or Site	Method	Annual Average	24-Hour Avg		Valid Data Recovery *	
			Max	2nd High	No. of Obs.	%
<b>Santa Cruz County</b>						
Nogales Post Office <sup>2</sup> (1)	FRM	13.1	49.7	33.0	60	98
Nogales Post Office <sup>2</sup> (2)	FRM	12.9	44.6	32.8	60	98

\***Valid data recovery** shows the number of valid observations during 2005 and the percentage of scheduled samples that were valid.

<sup>++</sup> Non Reference method.

<sup>1</sup> Samples collected every day – 365 sample days in 2005.

<sup>2</sup> Samples collected every sixth day - 61 sample days in 2005.

<sup>3</sup> Samples collected every third day - 122 sample days in 2005.

<sup>4</sup> Samples collected every hour - 8760 sample hours in 2005.

(1) Indicates the Primary monitor (used for NAAQS compliance) in a collocated pair of monitors.

(2) Indicates the Secondary monitor (used for precision and accuracy) in a collocated pair of monitors.

## Criteria Pollutants - Compliance

### Carbon Monoxide

There are two NAAQS for CO: an eight-hour standard (most critical for compliance) and a one-hour standard. The eight-hour standard is 9 ppm and the one-hour standard is 35 ppm. According to the Code of Federal Regulations, compliance for both standards is determined by having no more than one exceedance per calendar year. EPA determines attainment of the standard at all sites in the non-attainment (or monitoring) area by evaluating two calendar years of data from each site. The highest of the second-highest values in a two-year period must not exceed the standard of 9 ppm (greater than or equal to 9.5 ppm to adjust for rounding) for the eight-hour standard or 35 ppm (greater than or equal to 35.5 ppm) for the one-hour standard.

No exceedances of the one-hour or eight-hour standards were recorded in 2004 or 2005. The data are presented in Table 12 and Table 13.

*Table 12: 2004-2005 One-Hour Carbon Monoxide Compliance (in ppm)*

*NAAQS for one-hour carbon monoxide: The highest of the second-highest values in a two-year period must not exceed 35 ppm.*

*NOTE: Pinal County monitors closed in 2002.*

2004-2005 One-Hour Carbon Monoxide NAAQS Compliance Values by County		
County	Exceedances	Violations
Maricopa	0	0
Pima	0	0
<i>Summary: 20 of 20 monitors in compliance</i>		

**Table 12: 2004-2005 One-Hour Carbon Monoxide Compliance (in ppm)**

City or Site	2004		2005		Compliance Value
	Max Value	2nd High	Max Value	2nd High	
<b>Maricopa County</b>					
Buckeye <sup>S</sup>	0.9	0.9	1.1	1.1	1.1
Central Phoenix	5.0	4.4	5.2	5.1	5.1
Dysart <sup>S</sup>	2.1	1.8	1.7	1.7	1.8
Glendale <sup>S</sup>	6.1	3.2	3.2	3.1	3.2
Greenwood	7.6	7.3	5.9	5.4	7.3

<b>Table 12: 2004-2005 One-Hour Carbon Monoxide Compliance (in ppm)</b>					
<b>City or Site</b>	<b>2004</b>		<b>2005</b>		<b>Compliance Value</b>
	<b>Max Value</b>	<b>2nd High</b>	<b>Max Value</b>	<b>2nd High</b>	
JLG Supersite	4.9	4.9	5.6	5.1	5.1
Mesa <sup>S</sup>	3.0	2.6	3.4	3.3	3.3
North Phoenix <sup>S</sup>	4.1	3.7	3.8	3.5	3.7
South Phoenix <sup>S</sup>	6.7	5.9	5.5	5.2	5.9
South Scottsdale <sup>S</sup>	3.4	3.1	3.2	3.1	3.1
Tempe <sup>S</sup>	3.1	2.6	3.2	3.0	3.0
West Chandler <sup>S</sup>	2.9	2.7	3.5	2.7	2.7
West Indian School	6.9	6.7	6.8	6.5	6.7
West Phoenix	7.7	7.5	7.2	7.0	7.5
<b>Pima County</b>					
22nd St. & Alvernon	4.0	4.0	4.1	3.6	4.0
22nd St. & Craycroft	3.6	3.4	3.5	3.3	3.4
Cherry & Glenn <sup>S</sup>	4.0	3.9	3.8	3.4	3.9
Children's Park	2.2	2.2	2.0	1.8	2.2
Golf Links & Kolb <sup>S</sup>	3.6	3.5	3.3	3.2	3.5
Downtown	5.5	4.7	3.0	2.8	4.7

<sup>S</sup> Seasonal monitor. Maricopa County monitors operate during January 1 to April 1 and September 1 to December 31; 5088 hours in 2005. Pima County monitors operate during January 1 to May 1 and October 1 to December 31; 5118 hours in 2005.

*Table 13. 2004-2005  
Eight-Hour Carbon Monoxide  
Compliance (in ppm)*

*NAAQS for eight-hour carbon monoxide:  
The highest of the second-highest values in a  
two-year period must not exceed 9 ppm.  
NOTE: Pinal County monitors closed in 2002.*

2004-2005 Eight-Hour Carbon Monoxide NAAQS Compliance Values by County		
County	Exceedances	Violations
Maricopa	0	0
Pima	0	0
<i>Summary: 20 of 20 monitors in compliance</i>		

<b>Table 13: 2004-2005 Eight-Hour Carbon Monoxide Compliance (in ppm)</b>					
City or Site	2004		2005		Compliance Value
	Max Value	2nd High	Max Value	2nd High	
<b>Maricopa County</b>					
Buckeye <sup>S</sup>	0.5	0.4	0.9	0.9	0.9
Central Phoenix	3.4	3.3	4.1	3.8	3.8
Dysart <sup>S</sup>	1.1	1.1	1.3	1.2	1.2
Glendale <sup>S</sup>	2.4	2.1	2.4	2.3	2.3
Greenwood	4.9	4.3	4.2	4.1	4.3
JLG Supersite	4.2	4.0	3.7	3.6	4.0
Mesa <sup>S</sup>	1.7	1.7	2.4	2.4	2.4
North Phoenix <sup>S</sup>	2.2	2.0	2.3	2.2	2.2
South Phoenix <sup>S</sup>	3.5	3.3	3.8	3.2	3.3
South Scottsdale <sup>S</sup>	2.4	2.4	2.4	2.4	2.4
Tempe <sup>S</sup>	1.9	1.7	2.6	2.4	2.4
West Chandler <sup>S</sup>	2.1	2.1	2.4	2.0	2.1
West Indian School	4.7	4.6	5.3	4.8	4.8
West Phoenix	5.2	5.1	5.8	4.6	5.1
<b>Pima County</b>					
22nd St. & Alvernon	2.1	2.0	2.2	2.1	2.1
22nd St. & Craycroft	1.6	1.6	1.7	1.5	1.6
Cherry & Glenn <sup>S</sup>	2.7	2.2	2.5	2.4	2.4

City or Site	2004		2005		Compliance Value
	Max Value	2nd High	Max Value	2nd High	
Children's Park	1.4	1.4	1.1	1.1	1.4
Golf Links & Kolb <sup>S</sup>	2.1	2.1	2.2	2.1	2.1
Tucson Downtown	3.7	2.5	1.9	1.7	2.5

<sup>S</sup> Seasonal monitor. Maricopa County monitors operate during January 1 to April 1 and September 1 to December 31; 5088 hours in 2005. Pima County monitors operate during January 1 to May 1 and October 1 to December 31; 5112 hours in 2005.

### Nitrogen Dioxide

The NAAQS for NO<sub>2</sub> is 0.053 parts per million (ppm) for an annual average. The standard is attained when the annual arithmetic mean concentration in a calendar year is less than or equal to 0.053 ppm. To demonstrate attainment, the annual mean must be based upon hourly data that are at least 75 percent complete. NO<sub>2</sub> annual averages near Arizona power plants range from 2 percent to 17 percent of the standard; in the urban areas, from 30 percent to 70 percent. All Arizona sites were in compliance with the NAAQS. Refer to Table 6 for the 2005 averages.

County	Exceedances	Violations
La Paz	0	0
Maricopa	0	0
Pima	0	0
Pinal	0	0
<i>Summary: 11 of 11 monitors in compliance</i>		

### Sulfur Dioxide

There are three NAAQS for SO<sub>2</sub>, two primary (annual average and 24-hour block average) and one secondary (three-hour block average). The annual average standard is 0.030 ppm (80 µg/m<sup>3</sup>) and cannot be exceeded in a calendar year. The 24-hour block average standard is 0.14 ppm (365 µg/m<sup>3</sup>), not to be exceeded more than once per calendar year. A 24-hour block average is considered valid if at least 75 percent of the hourly averages for the 24-hour period are available. The 24-hour averages are determined from successive non-overlapping 24-hour blocks which begin at midnight each day. To demonstrate attainment, the second highest 24-hour block average must be based on hourly data that are at least 75 percent complete in each calendar quarter. A 24-hour block average is considered valid if 18 or more

valid hourly averages are available. The sum of the valid averages is divided by the number of valid hours to determine the 24-hour average.

The secondary three-hour standard is 0.5 ppm (1300  $\mu\text{g}/\text{m}^3$ ), not to be exceeded more than once per calendar year. The three-hour averages are determined from successive non-overlapping three-hour blocks starting at midnight each calendar day. To demonstrate attainment, the second highest three-hour average must be based upon hourly data that are at least 75 percent complete in each calendar quarter. All three hours of the block must be available to calculate a valid average. However, if only one or two hourly averages are available and the three-hour average would exceed the level of the standard when zeroes are substituted for the missing hours, the block would be considered valid.

In Arizona in 2005, the maximum concentration sites - all near copper smelters - comply with these standards; the concentrations ranging from 1 to 67 percent of the three-hour, 2 to 58 percent of the 24-hour and 42 percent of the annual average standards. Sites near power plants are close to background levels, with annual averages near 1  $\mu\text{g}/\text{m}^3$ . See Table 7 for the 2005 averages.

County	Annual		Three Hour		24-Hour	
	Exceedances	Violations	Exceedances	Violations	Exceedances	Violations
Gila	0	0	0	0	0	0
Maricopa	0	0	0	0	0	0
Pima	0	0	0	0	0	0
Pinal	0	0	0	0	0	0
<i>Summary: 15 out of 15 monitors in compliance</i>						

### *Ozone -- One-hour*

The NAAQS one-hour standard for ozone is 0.12 ppm. Compliance with this standard is attained when, for a three-year period, the expected number of days per calendar year with maximum hourly average concentrations above 0.12 ppm (0.124 ppm for rounding) is equal to or less than one. An exceedance day is defined as any day having one or more hourly averages equal to or greater than 0.125 ppm. Hourly averages for at least 75 percent of the hours sampled (18-24 hours per day) must be present. Only one exceedance of the one hour standard occurred in Arizona in 2005, a 0.129 ppm reading at Fountain Hills, in metropolitan Phoenix.

As there have been no violations of the one-hour O<sub>3</sub> standard since 1996, on May 15, 2001, EPA found that Maricopa County had reached attainment for the one-hour O<sub>3</sub> standard. A maintenance plan and redesignation request developed by Maricopa Association of Governments (MAG), demonstrating how the area will maintain compliance with the one-hour standard, was submitted to EPA on April 21, 2004.

### *Ozone -- Eight-hour*

On April 15, 2004, the Phoenix area was designated nonattainment for the new, more stringent, eight-hour ozone standard. The one-hour standard will be revoked one year following the effective date of the eight-hour designation or June 15, 2005. However, certain of the control measures developed and implemented for the one-hour standard are required to remain in place to ensure continued progress toward attainment of the new eight-hour standard.

EPA developed the eight-hour O<sub>3</sub> standard in response to human exposure studies that showed adverse health effects occur at lower ozone concentrations extending over several hours. After its proposal in 1997 and after a protracted legal battle, the eight-hour standard was officially promulgated in 2003 and nonattainment area boundaries established. The eight-hour ozone standard is 0.08 ppm (0.084 for rounding) for a daily maximum eight-hour average. This standard is met when the three-year average of the annual fourth-highest daily maximum eight-hour average O<sub>3</sub> concentration is less than or equal to 0.08 ppm. The data in Table 16 are for those sites in operation in 2003 – 2005.

**Table 16: 2003 to 2005  
Eight-Hour Ozone  
Compliance (in ppm)**

*NAAQS: The three-year average of the annual fourth-highest daily maximum eight-hour average ozone concentration is less than or equal to 0.08 ppm.*

2002 to 2005 Eight-Hour Ozone NAAQS Compliance Values, By County				
County	Eight-Hour Exceedances *			Sites in Violation
	2003	2004	2005	
Cochise	0	0	0	0
Coconino	0	0	0	0
Gila	3	0	0	0
Maricopa	32	1	3	0
Navajo	0	0	0	0
Pima	1	0	0	0
Pinal	5	0	0	0
Yavapai	0	0	0	0
Yuma	1	0	0	0
<i>Summary: 41 of 41 monitors in compliance for 2003 to 2005</i>				

\* Includes all eight-hour exceedances above fourth highest value.

**Table 16: 2003 to 2005 Eight-Hour Ozone Compliance (in ppm)**

**Bold** denotes exceedances and sites in violation.

City or Site	Fourth-Highest Value			Three-Year Average
	2003	2004	2005	
<b>Cochise County</b>				
Chiricahua NM Entrance Station	0.073	0.070	.072	0.072
<b>Coconino County</b>				
Grand Canyon NP – Hance	0.073	0.072	.079	0.075
<b>Gila County</b>				
Tonto NM <sup>S</sup>	0.084	0.077	.084	0.082
<b>La Paz County</b>				
Alamo Lake (Opened 05/20/05)	N/A	N/A	.075	N/A
<b>Maricopa County</b>				
Blue Point	<b>0.086</b>	0.075	.081	0.081

**Table 16: 2003 to 2005 Eight-Hour Ozone Compliance (in ppm)****Bold** denotes exceedances and sites in violation.

City or Site	Fourth-Highest Value			Three-Year Average
	2003	2004	2005	
Buckeye <sup>S</sup> (Opened 08/01/04)	N/A	0.058 #	.065	N/A
Cave Creek <sup>S</sup>	0.083	0.076	.082	0.080
Central Phoenix	0.079	0.074	.075	0.076
Dysart <sup>S</sup>	0.073	0.065	.066	0.068
Falcon Field <sup>S</sup>	0.079	0.070	.076	0.075
Fountain Hills	0.083	0.075	<b>.088</b>	0.082
Glendale <sup>S</sup>	<b>0.085</b>	0.076	.076	0.079
Humboldt Mt. <sup>S</sup>	<b>0.087</b>	0.078	<b>.087</b>	0.084
JLG Supersite	0.075	0.072	.076	0.074
North Phoenix	<b>0.086</b>	0.080	.084	0.083
Palo Verde <sup>S</sup> (Closed 10/31/2004)	0.075	0.072	N/A	N/A
Pinnacle Peak	0.083	0.068	.083	0.082
Rio Verde <sup>S</sup>	0.083	0.074	<b>.087</b>	0.081
South Phoenix	0.076	0.072	.076	0.075
South Scottsdale	0.079	0.073	.077	0.076
Tempe <sup>S</sup>	0.080	0.072	.076	0.076
West Chandler <sup>S</sup>	0.078	0.070	.075	0.074
West Phoenix	0.077	0.072	.068	0.072
<b>Navajo County</b>				
Petrified Forest NP	0.074	0.071	.070	.072
<b>Pima County</b>				
22nd St. & Craycroft	0.073	0.069	.074	0.072
Children's Park	0.076	0.068	.075	0.073
Coachline	0.061 #	0.068	.066	N/A
Green Valley	0.068 #	0.066	.068	N/A

**Table 16: 2003 to 2005 Eight-Hour Ozone Compliance (in ppm)**

**Bold** denotes exceedances and sites in violation.

City or Site	Fourth-Highest Value			Three-Year Average
	2003	2004	2005	
Rose Elementary	0.065 #	0.064	.067	N/A
Saguaro NP East	0.078	0.073	.079	0.077
Tangerine	0.074	0.068	.073	0.072
Tucson Downtown	0.068	0.063	.070	0.067
Tucson Fairgrounds	0.070	0.064	.073	0.069
<b>Pinal County</b>				
Apache Junction - Maintenance Yard	0.072	0.069	.068	0.069
Casa Grande - Airport	0.073	0.070	.072	0.072
Queen Creek <sup>S##</sup>	0.072	0.059	.067	0.066
Maricopa <sup>S</sup>	0.075	0.064	.061	0.066
Pinal Air Park <sup>S</sup>	0.074	0.067	.077	0.073
Queen Valley	0.087	0.073	.084	0.081
<b>Yavapai County</b>				
Hillside <sup>S</sup>	0.067	0.077	.074	0.073
<b>Yuma County</b>				
Yuma Game & Fish	0.078	0.073	.078	0.076

<sup>S</sup>Seasonal monitor, operational during April 1 to Nov. 1.

<sup>#</sup>Indicates the data do not satisfy EPA's summary criteria, usually meaning less than 75 percent valid data recovery available.

<sup>##</sup> Formerly "Combs"

N/A - Data not available

Notes: Data follow EPA truncation and averaging rules. Data published in previous annual reports may be slightly different.

### Particulate Matter - PM<sub>10</sub>

The NAAQS for particulate matter 10 microns and smaller in diameter (PM<sub>10</sub>) are 50 µg/m<sup>3</sup> for the annual arithmetic mean concentration and 150 µg/m<sup>3</sup> for the 24-hour average concentration.

The annual standard is attained when, for a three-year period, the expected annual arithmetic mean concentration is less than or equal to  $50\mu\text{g}/\text{m}^3$ . Annual arithmetic means are determined by calculating quarterly (three month) averages of the samples collected during that quarter; a minimum of 75 percent of the samples must be valid to produce the annual mean. This mean is rounded to the nearest  $1\mu\text{g}/\text{m}^3$  for comparison to the standard.

Compliance with the 24-hour  $\text{PM}_{10}$  standard is attained when the expected exceedance rate is one or less per year measured over three years. A sample value is rounded to the nearest  $10\mu\text{g}/\text{m}^3$  for comparison with the standard to determine if it is an exceedance (i.e., a sample value of  $154\mu\text{g}/\text{m}^3$  is not an exceedance; a sample value of  $155\mu\text{g}/\text{m}^3$  is an exceedance). Since the majority of monitoring sites do not collect daily samples, the expected exceedance rate must be calculated by quarter following EPA guidelines. The same requirements of 75 percent completeness and three consecutive years of data apply.

Tables 17 and 18 present the 2003 to 2005 expected exceedance rates for the  $\text{PM}_{10}$  annual arithmetic means and maximum 24-hour average values

*Table 17: 2003 to 2005 Annual Average  $\text{PM}_{10}$  Compliance (in  $\mu\text{g}/\text{m}^3$ , Standard Conditions)*

*NAAQS: The expected annual arithmetic mean (average of three most recent annual means) is less than or equal to  $50\mu\text{g}/\text{m}^3$ .*

*The expected annual arithmetic mean is rounded to the nearest  $1\mu\text{g}/\text{m}^3$  for comparison to the standard.*

2003 to 2005 $\text{PM}_{10}$ Annual Average NAAQS Compliance Values, By County				
County	Sites above Standard			Sites in Violation
	2003	2004	2005	
Apache	0	0	0	0
Cochise	0	0	0	0
Coconino	0	0	0	0
Gila	0	0	0	0
Graham	0	0	0	0
Maricopa	5	2	4	4
Mohave	0	0	0	0
Navajo	0	0	0	0
Pima	0	0	0	0
Pinal	1	1	2	1
Santa Cruz	0	0	1	0
Yavapai	0	0	0	0
Yuma	0	0	0	0
<i>Summary: 54 of 59 monitors in compliance</i>				

<b>Table 17: 2003 to 2005 Annual Average PM<sub>10</sub> Compliance (in µg/m<sup>3</sup>)</b>				
<b>City or Site</b>	<b>2003</b>	<b>2004</b>	<b>2005</b>	<b>Expected Annual Mean</b>
<b>Apache County</b>				
TEP – Springerville – Coalyard	16.2	13.4	15.4	15
TEP – Springerville – Coyote Hills	11.9	10.2	10.3	11
<b>Cochise County</b>				
Douglas – Red Cross	30.3	26.3	34.4	30
Paul Spur	19.3	14.7	27.6 #	N/A
<b>Coconino County</b>				
Flagstaff – Middle School	19.8 #	16.0	17.0 #	N/A
Sedona	26.5 #	11.1	12.2 #	N/A
<b>Gila County</b>				
Hayden – Old Jail	36.0	27.5	29.9 #	N/A
Miami – Golf Course	21.0	16.9	21.0	20
Miami – Ridgeline	15.0	10.2	12.4	13
Payson	24.3	18.9	22.1#	N/A
<b>Graham County</b>				
Safford	23.4	17.0	20.8#	N/A
<b>Maricopa County</b>				
Bethune Elementary School	47.2	42.4	58.6	49
Buckeye (Opened 10/19/2004)	N/A	39.9 #	52.7	N/A
Central Phoenix – every 6th day monitor	39.8	32.3	38.5	37
Central Phoenix – continuous monitor	43.0	36.6	37.1	39
Chandler	49.8	39.6	49.4	46
Durango Complex	<b>62.3</b>	<b>51.6</b>	<b>66.4</b>	<b>60</b>

<b>Table 17: 2003 to 2005 Annual Average PM<sub>10</sub> Compliance (in µg/m<sup>3</sup>)</b>				
<b>City or Site</b>	<b>2003</b>	<b>2004</b>	<b>2005</b>	<b>Expected Annual Mean</b>
Dysart (Opened 7/16/2003)	36.3 #	27.3	29.0	N/A
Glendale	35.6	25.7	29.0	30
Greenwood	<b>51.0</b>	44.3	<b>52.3</b>	49
Higley	<b>61.6</b>	47.9	<b>51.4</b>	<b>54</b>
JLG Supersite (Closed 12/31/2003 – part of urban haze program; Reopened 1/1/2005)	36.6 #	N/A	32.3	N/A
Mesa	33.6	23.2	30.0	29
North Phoenix	33.9	24.8	29.6	29
Palo Verde (Closed 12/31/2004)	26.4	14.5	N/A	N/A
South Phoenix	<b>52.0</b>	45.6	<b>54.7</b>	<b>51</b>
South Scottsdale	36.1	26.1	34.0	32
West Chandler	42.4	29.9	34.2	36
West Forty Third	<b>62.3</b>	<b>61.1</b>	<b>73.9</b>	<b>66</b>
West Phoenix	46.4	36.9	44.5	43
<b>Mohave County</b>				
Bullhead City – ADEQ	20.1	18.2	18.6 #	N/A
<b>Navajo County</b>				
Show Low	18.2	14.9	13.7 #	N/A
<b>Pima County</b>				
Ajo	22.7	19.3	22.7	22
Broadway & Swan	26.6	20.7	23.7	24
Corona de Tucson	16.8	12.4	15.4	15
Green Valley	18.7	13.6	17.4	17
Orange Grove	29.8	26.8	29.2	29
Prince Road	31.4	28.4	37.0 #	30

<b>Table 17: 2003 to 2005 Annual Average PM<sub>10</sub> Compliance (in µg/m<sup>3</sup>)</b>				
<b>City or Site</b>	<b>2003</b>	<b>2004</b>	<b>2005</b>	<b>Expected Annual Mean</b>
Rillito, ADEQ	39.5	32.2	39.1	37
Rillito, APCC	33.6	26.9	26.8	29
Santa Clara	26.7	20.4	26.5	25
South Tucson	34.1	29.2	30.2	31
Tangerine	19.3	14.7	19.1	18
<b>Pinal County</b>				
Apache Junction Fire Station (Opened 7/1/2003)	26.7 #	18.4	19.9	N/A
Casa Grande Downtown	31.5	24.4	30.9	29
Coolidge Maintenance Yard	35.3	24.5	36.0	32
Eloy	41.5	27.8	33.4	34
Mammoth	16.4	11.8	13.6	14
Pinal Air Park	28.6	20.2	22.3	24
Pinal County Housing Complex	<b>61.0</b>	47.1	<b>56.7 #</b>	<b>N/A</b>
Riverside Maintenance Yard (Opened 3/2003)	23.9 #	15.2	18.1	N/A
Stanfield	46.1 #	33.9	<b>52.2</b>	N/A
<b>Santa Cruz County</b>				
Nogales Post Office	37.5	42.6	<b>56.9</b>	46
<b>Yavapai County</b>				
Clarkdale – NW (#2)	19.4	14.7	14.7	16.3
Clarkdale – SE (#1)	23.8	19.8	21.8	21.8
Prescott Valley (Opened 3/12/2003)	13.9 #	12.9	14.8 #	N/A
<b>Yuma County</b>				
Yuma – Juvenile Center/Courthouse	38.1	35.5 #	34.9	N/A

<b>Table 17: 2003 to 2005 Annual Average PM<sub>10</sub> Compliance (in µg/m<sup>3</sup>)</b>				
<b>City or Site</b>	<b>2003</b>	<b>2004</b>	<b>2005</b>	<b>Expected Annual Mean</b>
<b>Mexico</b>				
Agua Prieta – Fire Station	<b>60.3</b>	<b>60.5</b>	<b>68.1</b>	<b>63</b>
Nogales – Fire Station	<b>65.0 #</b>	50.2	<b>62.9</b>	<b>N/A</b>

Bold denotes value above the standard.

N/A – Not available

# Indicates the data do not satisfy EPA's summary criteria, usually meaning less than 75 percent valid data recovery available in one or more calendar quarters.

Notes:

For collocated sites, data from the Primary monitor (POC 1) are used for the Annual Average calculations. However, if valid data recovery is between 50% and 75%, data from the Secondary (POC2) monitor can be used. If no Secondary data are available, data substitution can be made following the EPA document, 'Guideline on Exceptions to Data Requirements for Determining Attainment of Particulate Matter Standards.'

**Table 18: 2003 to 2005 Maximum 24-Hour Average PM<sub>10</sub> Compliance (in µg/m<sup>3</sup>, Standard Conditions)**

*NAAQS: Expected occurrence of exceedances (samples equal to or greater than 150 µg/m<sup>3</sup>) is one or less over three consecutive years.*

*Sample values are rounded to the nearest 10 µg/m<sup>3</sup> to determine exceedance; values less than or equal to 154 µg/m<sup>3</sup> are not exceedances; values greater than or equal to 155 µg/m<sup>3</sup> are exceedances.*

2003 to 2005 PM <sub>10</sub> Maximum 24-Hour Compliance Values, By County				
	Sites with Exceedances			Sites in Violation
	2003	2004	2005	
Apache	0	0	0	0
Cochise	1	0	0	1
Coconino	0	0	0	0
Gila	0	0	0	0
Graham	0	0	0	0
Maricopa	14	1	6	12
Mohave	0	0	0	0
Navajo	0	0	0	0
Pima	1	0	0	0
Pinal	2	2	2	2
Santa Cruz	1	0	1	1
Yavapai	0	0	0	0
Yuma	0	0	0	0
<i>Summary: 42 of 58 monitors in compliance</i>				

**Table 18: 2003 to 2005 Maximum 24-Hour Average PM<sub>10</sub> Compliance (in µg/m<sup>3</sup>)**

City or Site	2003		2004		2005		3-Year Avg Expected Rate of Exceedance
	Max 24-Hr Avg	Exp. Exc.	Max 24-Hr Avg	Exp. Exc.	Max 24-Hr Avg	Exp. Exc.	
<b>Apache County</b>							
TEP – Springerville – Coalyard	185.0	1	128.7	0	197.6	1	<1
TEP – Springerville – Coyote Hills	76.8	0	68.6	0	28.9	0	<1
<b>Cochise County</b>							
Douglas Red Cross	79	0	56	0	86	0	<1
Paul Spur Chemical Lime Plant	<b>207</b>	<b>6.4</b>	44	0	76	0	<b>2.1</b>
<b>Coconino County</b>							
Flagstaff Middle School	60 #	0	42	0	38	0	<1

<i>Table 18: 2003 to 2005 Maximum 24-Hour Average PM<sub>10</sub> Compliance (in µg/m<sup>3</sup>)</i>							
City or Site	2003		2004		2005		3-Year Avg Expected Rate of Exceedance
	Max 24-Hr Avg	Exp. Exc.	Max 24-Hr Avg	Exp. Exc.	Max 24-Hr Avg	Exp. Exc.	
Sedona Post Office	69 #	0	32	0	34	0	<1
<b>Gila County</b>							
Hayden – Old Jail	91	0	55	0	124	0	<1
PDMI - Miami – Golf Course	53	0	40	0	40	0	<1
PDMI - Miami – Ridgeline	59	0	26	0	23	0	<1
Payson Well Site	99	0	52	0	81	0	<1
<b>Graham County</b>							
Safford	76	0	99	0	50	0	<1
<b>Maricopa County</b>							
Bethune Elementary School (Opened 1/3/2003)	145	0	122	0	<b>198</b>	<b>6.4</b>	<b>2.1</b>
Buckeye <sup>E</sup> (Opened 10/19/2004)	N/A	N/A	82 #	0	<b>169</b>	<b>2</b>	N/A
Central Phoenix – every 6th day monitor	114	0	81	0	125	0	<1
Central Phoenix – continuous monitor <sup>E</sup>	<b>183</b>	<b>3.1</b>	94	0	116	0	1.0
Chandler	<b>240</b>	<b>6.0</b>	150	0	130	0	<b>2.0</b>
Durango Complex <sup>E</sup>	<b>195</b>	<b>6.0</b>	139	0	<b>206</b>	<b>13.0</b>	<b>6.3</b>
Dysart (Opened 7/16/2003)	133 #	0	94	0	76	0	<1
Glendale	151	0	69	0	84	0	<1
Greenwood	<b>166</b>	<b>6.0</b>	100	0	<b>173</b>	<b>6</b>	<b>4.0</b>
Higley <sup>E</sup>	<b>225</b>	<b>6.0</b>	<b>159</b>	<b>1</b>	142	0	<b>2.3</b>

<b>Table 18: 2003 to 2005 Maximum 24-Hour Average PM<sub>10</sub> Compliance (in µg/m<sup>3</sup>)</b>							
<b>City or Site</b>	<b>2003</b>		<b>2004</b>		<b>2005</b>		<b>3-Year Avg Expected Rate of Exceedance</b>
	<b>Max 24-Hr Avg</b>	<b>Exp. Exc.</b>	<b>Max 24-Hr Avg</b>	<b>Exp. Exc.</b>	<b>Max 24-Hr Avg</b>	<b>Exp. Exc.</b>	
JLG Supersite (Closed 12/31/2003 – part of urban haze program)	169 #	6.0	N/A	N/A	N/A	N/A	N/A
Maryvale (Closed 4/1/2004)	151	0	46 #	0	N/A	0	N/A
Mesa	176	6.0	49	0	86	0	2.0
North Phoenix	155	6.0	46	0	81	0	2.0
Palo Verde (Closed 01/05/2005)	158	6.4	42	0	N/A	0	N/A
South Phoenix	164	6.0	132	0	147	0	2.0
South Scottsdale	172	6.0	77	0	121	0	2.0
West Chandler <sup>E</sup>	206	13.7	70	0	94	0	4.6
West Forty Third <sup>E</sup>	157	6.0	145	0	233	13.0	6.3
West Phoenix	158	6.4	100	0	155	6.0	4.1
<b>Mohave County</b>							
Bullhead City – ADEQ	121	0	48	0	48	0	<1
<b>Navajo County</b>							
Show Low	58	0	41	0	37	0	<1
<b>Pima County</b>							
Ajo – ADOT	139	0	43	0	45	0	<1
Broadway & Swan	122	0	35	0	46	0	<1
Corona De Tucson	104	0	37	0	33	0	<1
Green Valley	127	0	127	0	54	0	<1
Orange Grove <sup>E</sup>	152	0	119	0	98	0	<1
Prince Road	126	0	67	0	88	0	<1
Rillito , ADEQ	118	0	93	0	84	0	<1

<b>Table 18: 2003 to 2005 Maximum 24-Hour Average PM<sub>10</sub> Compliance (in µg/m<sup>3</sup>)</b>							
<b>City or Site</b>	<b>2003</b>		<b>2004</b>		<b>2005</b>		<b>3-Year Avg Expected Rate of Exceedance</b>
	<b>Max 24-Hr Avg</b>	<b>Exp. Exc.</b>	<b>Max 24-Hr Avg</b>	<b>Exp. Exc.</b>	<b>Max 24-Hr Avg</b>	<b>Exp. Exc.</b>	
Rillito , APCC (1-in-3 day schedule)	<b>256</b>	<b>3.1</b>	130	0	83.7	0	1
Santa Clara	146	0	41	0	82	0	<1
South Tucson	150	0	149	0	73	0	<1
Tangerine	125	0	34	0	37	0	<1
<b>Pinal County</b>							
Apache Junction Maintenance Yard (Closed 12/31/2003)	95	0	N/A	N/A	N/A	N/A	N/A
Apache Junction Fire Station (Opened 7/1/2003)	103	0	35	0	47	0	N/A
Casa Grande Downtown	99	0	52	0	79	0	<1
Coolidge Maintenance Yard	106	0	57	0	81	0	<1
Eloy City Complex	154	0	46	0	73	0	<1
Mammoth County Complex	89	0	30	0	33	0	<1
Pinal Air Park	108	0	39	0	122	0	<1
Pinal County Housing Complex	<b>289</b>	<b>12.0</b>	<b>155</b>	<b>6</b>	<b>179</b>	<b>12.0</b>	<b>10.0</b>
Riverside Maintenance Yard (Opened 3/2003)	101 #	0	34	0	35	0	N/A
Stanfield	<b>171 #</b>	<b>6.4</b>	80	0	<b>173</b>	<b>6</b>	<b>4.1</b>
<b>Santa Cruz County</b>							
Nogales Post Office	<b>184</b>	<b>12.3</b>	<b>140</b>	<b>0</b>	<b>280</b>	<b>12</b>	<b>8.1</b>
<b>Yavapai County</b>							
Clarkdale – NW (#2)	68	0	36	0	31.5	0	0
Clarkdale – SE (#1)	59	0	41	0	43.1	0	0

<b>Table 18: 2003 to 2005 Maximum 24-Hour Average PM<sub>10</sub> Compliance (in µg/m<sup>3</sup>)</b>							
<b>City or Site</b>	<b>2003</b>		<b>2004</b>		<b>2005</b>		<b>3-Year Avg Expected Rate of Exceedance</b>
	<b>Max 24-Hr Avg</b>	<b>Exp. Exc.</b>	<b>Max 24-Hr Avg</b>	<b>Exp. Exc.</b>	<b>Max 24-Hr Avg</b>	<b>Exp. Exc.</b>	
Prescott Valley (Opened 3/12/2003)	68 #	0	31	0	53	0	N/A
<b>Yuma County</b>							
Yuma – Juvenile Center/Courthouse	127	0	114 #	0	94	0	0

Bold denotes value above the standard.

N/A – Not available

# Indicates the data do not satisfy EPA’s summary criteria, usually meaning less than 75 percent valid data recovery available in one or more calendar quarters.

<sup>E</sup> Indicates every day monitoring. Phoenix area sites which began every day monitoring in 2004 include: Buckeye, Durango Complex, Higley, and West Forty Third.

### **Particulate Matter – PM<sub>2.5</sub>**

The NAAQS for particulate matter 2.5 microns and smaller in diameter (PM<sub>2.5</sub>) are 15.0 micrograms per cubic meter (µg/m<sup>3</sup>) for the annual arithmetic mean concentration and 65 µg/m<sup>3</sup> for the 24-hour average concentrations. Appendix N to Part 50 of the 40 CFR will be used to assess the compliance of the monitors operating in Arizona during 2005.

The annual PM<sub>2.5</sub> standard is met when the three-year average of annual means is less than or equal to 15.0 µg/m<sup>3</sup>. This three-year average is determined by calculating the quarterly averages for each year (with 75 percent data recovery in each quarter) to determine the calendar year average and then averaging the three years together.

The 24-hour standard is met when the three-year average of the 98th percentile values is less than or equal to 65 µg/m<sup>3</sup>. There must also be 75 percent data completeness for each year.

Please note that the data in Table 19 are from federal reference monitors. In prior years, the dichot fine measurement was used as an approximate equivalent for PM<sub>2.5</sub>, but the federal reference monitors provide a more accurate measurement of this pollutant. Data are collected and reported in local conditions.

In February of 2004, Arizona requested that all parts of the State (except for Indian Country) be designated attainment/unclassifiable for the PM<sub>2.5</sub> NAAQS.

*Table 19: 2003 to 2005 Annual Average PM<sub>2.5</sub> Compliance (in µg/m<sup>3</sup>, local conditions)*

*NAAQS: The three-year average of annual means is less than or equal to 15 µg/m<sup>3</sup>*

2003 to 2005 PM <sub>2.5</sub> Annual Average NAAQS Compliance Values, By County				
	Sites with Exceedances			Sites in Violation
	2003	2004	2005	
Cochise	0	0	0	0
Coconino	0	0	0	0
Gila	0	0	0	0
Maricopa	0	0	0	0
Pima	0	0	0	0
Santa Cruz	0	0	0	0
<i>Summary: 10 of 10 federal reference monitors in compliance</i>				

<b>Table 19: 2003 to 2005 Annual Average PM<sub>2.5</sub> Compliance (in µg/m<sup>3</sup>)</b>				
<b>City or Site Federal Reference Monitors</b>	<b>2003</b>	<b>2004</b>	<b>2005</b>	<b>Three- Year Average</b>
<b>Cochise County</b>				
Douglas Red Cross	6.47 #	7.11	7.33	N/A
<b>Coconino County</b>				
Flagstaff Middle School	5.69 #	6.77	6.01	N/A
<b>Gila County</b>				
Payson Well Site	9.01 #	9.54	8.38 #	N/A
<b>Maricopa County</b>				
JLG Supersite	11.27	9.73	9.72	10.2
Tempe Community Center (Closed 7/26/2004)	9.63	7.30 #	N/A	N/A
West Phoenix	10.68	11.60	11.87	11.4
<b>Pima County</b>				
Children's Park	6.54	5.57	5.91	6.0
Orange Grove	6.45	5.79	6.32	6.2

**Table 19: 2003 to 2005 Annual Average PM<sub>2.5</sub> Compliance (in µg/m<sup>3</sup>)**

City or Site Federal Reference Monitors	2003	2004	2005	Three- Year Average
<b>Pinal County</b>				
Apache Junction Fire Station	6.30 #	5.51 #	5.52	5.8
Casa Grande Downtown	8.42	7.13	7.33	7.62
<b>Santa Cruz County</b>				
Nogales Post Office	11.30	10.83	13.1	11.4

# Indicates the data do not satisfy EPA's summary criteria, usually meaning less than 75 percent valid data recovery available in one or more calendar quarters.

**Table 20: 2003 to 2005 24-Hour Average PM<sub>2.5</sub> Compliance (in µg/m<sup>3</sup>, local conditions)**

*NAAQS: The three-year average of the 98th percentile values is less than or equal to 65 µg/m<sup>3</sup>.*

*Note: The three-year average is rounded to the nearest 1 µg/m<sup>3</sup> for comparison to the standard.*

2003 to 2005 PM <sub>2.5</sub> 24-Hour Average NAAQS Compliance Values, By County				
	Sites with Exceedances			Sites in Violation
	2003	2004	2005	
Cochise	0	0	0	0
Coconino	0	0	0	0
Gila	0	0	0	0
Maricopa	0	0	0	0
Pima	0	0	0	0
Santa Cruz	0	0	0	0
<i>Summary: 10 of 10 federal reference monitors in compliance</i>				

<b>Table 20. 2003 to 2005 24-Hour Average PM<sub>2.5</sub> Compliance (in µg/m<sup>3</sup>)</b>				
<b>City or Site Federal Reference Monitors</b>	<b>98th Percentile Samples **</b>			<b>Three- Year Average</b>
	<b>2003</b>	<b>2004</b>	<b>2005</b>	
<b>Cochise County</b>				
Douglas Red Cross	11.7#	22.5	16.0	16.7
<b>Coconino County</b>				
Flagstaff Middle School	16.9 #	20.7	12.7	16.8
<b>Gila County</b>				
Payson	24.9 #	19.3	22.9 #	22.4
<b>Maricopa County</b>				
JLG Supersite	24.2	27.6	28.2	24.1
Tempe Community Center (Closed 7/26/2004)	25.0	14.8 #	N/A	N/A
West Phoenix	25.9	29.9	40.5	32.1
<b>Pima County</b>				
Children's Park	13.2	10.3	10.7	11.4
Orange Grove	15.9	13.3	13.7	14.3
<b>Pinal County</b>				
Apache Junction Fire Station	21.1	10.3	10.6	14.0
Casa Grande Downtown	26.7	13.7	16.9	19.1
<b>Santa Cruz County</b>				
Nogales – Post Office	35.0	25.1	33.0	31.0

\*\* The 98th percentile value will be the second highest value for sites on an every 6th day sample schedule. The 98th percentile value will be the 3rd highest value for sites on an every 3rd day sample schedule.

# Indicates the data do not satisfy EPA's summary criteria, usually meaning less than 75 percent valid data recovery available in one or more calendar quarters.

## Visibility Data

Visibility monitoring is of three types: aerosol, optical and scene. Aerosol measurements include the physical properties of the ambient atmospheric particles (chemical composition, size, shape, concentration, temporal and spatial distribution and other physical properties) through which a scene is viewed. The chemical species that comprise a particulate sample have different extinction efficiencies. Extinction efficiency is the extent to which an individual or a specific particle will either scatter or absorb light, thus blocking the light's path to one's eye. The overall impact of particles can be estimated by summing the effect of all the component species. This method is the primary approach used in the national regional haze rule for estimating present visibility and charting trends for future plan reviews. Optical methods measure either light scattering or light extinction continuously. Scene measurements are photograph-based with subsequent analysis.

ADEQ operates several types of monitors designed to characterize different optical phenomena. Visibility data from these monitors can be expressed by several different measurement units: deciview, inverse megameters, and visual range. Inverse megameters is a representation of the ratio between how much light is not received by a sensor compared to the amount of light that leaves a source. Higher numbers mean worse visibility.

## Class I Areas

In anticipation of the federal regional haze rule, ADEQ, in 1997, undertook development of a visibility monitoring program directed at Class I areas in partnership with Arizona's federal land managers. The aim is to collect data at all of Arizona's Class I areas. Based on the regional haze rule, five years of data will be needed to determine baseline and projected visibility conditions. Since the IMPROVE program consists of aerosol sampling only, ADEQ included nephelometers for measuring light scattering at its jointly operated sites. IMPROVE aerosol samplers operate every three days and represent 24-hour averages. Taking continuous measurements provides insight into variation in visibility impairment with time, along with advancing the understanding of the relationship between particles and light scattering.

Table 21 summarizes the nephelometer data from locations in or near Arizona Class I areas from 1998 to 2005. The data are summarized into three categories for all hours (24 hours a day): the average visibility of the dirtiest 20 percent of the sampled hours, the mean visibility of all hours and the average visibility of the cleanest 20 percent of the sampled hours. As natural background levels are  $15 \text{ Mm}^{-1}$ , this table shows that on average most sites are within background, with the exceptions being Tucson Mountain in 2002 – 2005, and Pleasant Valley in 2003.

<b>Table 21: Visibility in Class I Areas (Nephelometer Data in <math>Mm^{-1}</math>)</b>				
<b>Site and Wilderness Area</b>	<b>Year</b>	<b><math>Mm^{-1}</math> (24 hour Averages)</b>		
		<b>Mean of the 20% Dirtiest Sampled Hours</b>	<b>Mean of all Sampled Hours</b>	<b>Mean of the Cleanest 20% Sampled Hours</b>
Greer Water Treatment Plant <i>Mt. Baldy Wilderness</i>	2002	26	10	2
	2003	26	10	1
	2004	17	8	1
	2005	23	9	1
Humboldt Mountain <i>Mazatzal Wilderness and Pine Mountain Wilderness</i> <i>(Site closed in 2004)</i>	1998	24	9	0
	1999	25	12	3
	2000	28	13	3
	2001	21	9	1
	2002	24	8	0
	2003	36	16	3
Ike's Backbone <i>Mazatzal/Pine Mountain Wildernesses</i>	2002	24	10	2
	2003	30	12	2
	2004	24	11	3
	2005	26	12	4
Mount Ord <i>Mazatzal Wilderness (site closed in 2000)</i>	1998	28	12	2
	1999	22	11	3
McFadden Peak <i>Sierra Ancha Wilderness (site closed in 2000)</i>	1998	24	10	1
	1999	18	7	0
Muleshoe Ranch <i>Chiracahua National Monument Wilderness, Galiuro Wilderness, Chiricahua Forest</i>	1998	24	11	4
	1999	20	11	3
	2000	22	11	3
	2001	24	12	4

<b>Table 21: Visibility in Class I Areas (Nephelometer Data in <math>Mm^{-1}</math>)</b>				
<b>Site and Wilderness Area</b>	<b>Year</b>	<b><math>Mm^{-1}</math> (24 hour Averages)</b>		
		<b>Mean of the 20% Dirtiest Sampled Hours</b>	<b>Mean of all Sampled Hours</b>	<b>Mean of the Cleanest 20% Sampled Hours</b>
<i>Service Wilderness</i>	2002	25	12	4
	2003	25	11	3
	2004	20	8	1
	2005	21	10	4
Rucker Canyon <i>Chiricahua Wilderness</i> <i>(site closed in 2001)</i>	1998	30	12	3
	1999	20	10	4
	2000	18	8	1
Pleasant Valley Ranger Station <i>Sierra Ancha Wilderness</i>	2001	28	14	5
	2002	27	13	3
	2003	33	15	4
	2004	20	10	3
	2005	28	13	4
Camp Raymond <i>Sycamore Canyon Wilderness</i>	1998	N/A	N/A	N/A
	1999	28	13	4
	2000	28	13	3
	2001	28	13	3
	2002	30	13	3
	2003	32	14	3
	2004	25	12	3
	2005	33	14	3
Tucson Mountain <i>Saguaro National Park</i>	1998	30	12	2
	1999	24	13	6

**Table 21: Visibility in Class I Areas (Nephelometer Data in  $Mm^{-1}$ )**

Site and Wilderness Area	Year	$Mm^{-1}$ (24 hour Averages)		
		Mean of the 20% Dirtiest Sampled Hours	Mean of all Sampled Hours	Mean of the Cleanest 20% Sampled Hours
<i>(Includes both the West facilities support building and the National Park Service well site)</i>	2000	23	12	5
	2001	22	11	3
	2002	31	16	6
	2003	35	17	6
	2004	32	16	5
	2005	31	16	5
Chiricahua National Monument	2004	18	9	3
	2005	21	10	2
Organ Pipe National Monument	2004	21	10	3
	2005	23	12	4
Petrified Forest National Park	2004	20	9	3
	2005	24	11	3

*N/A – Not available*

### Urban Haze

Besides the Class I areas, ADEQ also operates transmissometers and nephelometers in Phoenix and Tucson. Data from these instruments through 2005 are presented in Table 22. The data are separated into categories for all hours and for 6-hours. Each category is further summarized into the average visibility for the dirtiest 20 percent of the sampled hours, the mean visibility of all hours and the cleanest 20 percent of the sampled hours. As visual range in miles may be a more familiar unit, the values in inverse megameters ( $Mm^{-1}$ ) in Table 22 can be converted to visual range in miles by the expression  $(2431/b_{ext})$ . A few conversions are given here:

<u><math>Mm^{-1}</math></u>	<u>Miles</u>	<u>Comment</u>
133	18	Highest in Table 22
100	24	
50	48	
7	347	Lowest in the Table

**Table 22. Phoenix and Tucson Urban Haze Data 1998 to 2005 (in  $Mm^{-1}$ )**

Site	Year	24 Hour Samples			5 a.m. to 11 a.m.		
		Dirtiest 20%	Mean	Cleanest 20%	Dirtiest 20%	Mean	Cleanest 20%
Mesa Transmissometer	2004	106	60	24	110	65	29
	2005	121	72	35	123	78	44
Phoenix Transmissometer	1998	133	78	45	136	84	50
	1999	127	72	38	128	77	42
	2000	131	74	38	134	80	42
	2001	118	69	36	118	73	42
	2002	124	75	42	125	79	46
	2003	131	72	36	135	78	42
	2004	121	69	35	126	75	42
	2005	126	72	36	128	78	43
Phoenix Nephelometer (Supersite)	1998	91	35	10	77	34	13
	1999	87	36	11	74	36	14
	2000	93	39	12	80	39	15
	2001	73	32	12	66	33	15
	2002	72	33	12	62	33	14
	2003	79	34	11	73	35	14
	2004	72	30	9	61	30	11
	2005	80	33	9	73	33	11

**Table 22. Phoenix and Tucson Urban Haze Data 1998 to 2005 (in  $Mm^{-1}$ )**

Site	Year	24 Hour Samples			5 a.m. to 11 a.m.		
		Dirtiest 20%	Mean	Cleanest 20%	Dirtiest 20%	Mean	Cleanest 20%
Phoenix Nephelometer (Dysart)	2004	46	22	7	52	27	9
	2005	41	20	8	41	23	10
Phoenix Nephelometer (Estrella Mountain)	2004	54	24	7	68	32	10
	2005	76	35	12	77	39	14
Phoenix Nephelometer (Vehicle Emissions)	2004	69	29	9	64	31	12
	2005	76	35	12	73	37	15
Tucson Transmissometer	1998	102	57	28	119	69	34
	1999	90	57	35	107	65	38
	2000	98	56	27	114	66	31
	2001	96	55	26	109	66	33
	2002	87	49	24	109	61	29
	2003	88	52	26	107	62	30
	2004	97	58	27	113	67	32
	2005	101	61	31	125	76	39
Tucson Nephelometer (U of A Central)	1998	45	21	4	47	23	7
	1999	43	23	10	41	24	11
	2000	40	20	8	40	22	9
	2001	42	23	10	44	25	13
	2002	38	20	7	42	22	9

<b>Table 22. Phoenix and Tucson Urban Haze Data 1998 to 2005 (in <math>Mm^{-1}</math>)</b>							
<b>Site</b>	<b>Year</b>	<b>24 Hour Samples</b>			<b>5 a.m. to 11 a.m.</b>		
		<b>Dirtiest 20%</b>	<b>Mean</b>	<b>Cleanest 20%</b>	<b>Dirtiest 20%</b>	<b>Mean</b>	<b>Cleanest 20%</b>
			2003	43	23	9	45
	2004	38	20	8	42	22	10
	2005	45	24	10	47	27	12
Tucson Nephelometer (Craycroft)	2001	38	19	8	N/A	N/A	N/A
	2002	37	18	7	N/A	N/A	N/A
	2003	52	25	7	N/A	N/A	N/A
	2004	42	21	8	43	22	9
	2005	35	19	7	44	25	11
Tucson Nephelometer (Children's Park)	2004	41	20	8	43	23	10
	2005	35	19	7	35	20	8

*N/A – Not available*

## Special Projects

---

### *Introduction*

In addition to ADEQ's statewide regulatory ambient air monitoring program, the Air Quality Division continued several special projects during 2005 and the first half of 2006. All of these studies go beyond data collection and seek to provide a better understanding of air pollutant science in Arizona and the Southwest. Data are employed in advanced computer models that help to explain and predict the relationship between emissions and air pollutant concentrations under a variety of conditions. Control strategies are modeled to predict the most effective methods to attain and maintain the National Ambient Air Quality Standards in Arizona. Issues related to the international border, identification of potential air pollution hotspots, improved visibility and reduction of regional haze, and appropriate responses to smoke and other air pollution hazards to protect public health fall under special projects. The knowledge gained from these studies can then be used by decision-makers to choose the most effective control strategies that will continue to improve the State's air quality.



**Figure 4 - Yuma West Monitoring Station, Western Arizona/Sonora Border Air Quality Study**

### *Yuma PM<sub>10</sub> Nonattainment Area Redesignation Project*

Yuma was designated nonattainment for PM<sub>10</sub> (particulate matter 10 microns or smaller) in 1990. ADEQ developed a State Implementation Plan (SIP) for Yuma in 1991 that demonstrated the area could meet the federal NAAQS by December 1994. After several consecutive years of clean monitoring data, a stakeholder process to prepare an attainment demonstration and maintenance plan was convened in July 2001. ADEQ met with local stakeholders to review the control measures already in place and hired a contractor to assist in developing an emissions inventory for the 1999 base year and future years emissions estimates. After air quality modeling for 1999 was completed successfully, ADEQ staff learned that incomplete monitoring data for 2001 would necessitate using the 2002-2004 monitoring data for the attainment demonstration, with a SIP submittal in early 2005.

On August 18, 2002, however, an unusually large and intense thunderstorm with blowing dust over east-central Sonora moved northwesterly through Yuma. For this day there were three hours with wind speeds above the dust re-suspension threshold of 15 mph. The Yuma PM<sub>10</sub> monitor registered 170 ug/m<sup>3</sup>, exceeding the National Ambient Air Quality Standard of 150 ug/m<sup>3</sup>. Data from nearby meteorological sites were tested to determine whether the exceedance date in question is considered meteorologically exceptional. These tests are described in an ADEQ document, “Technical Criteria Document for Determination of Natural Exceptional Events for Particulate Matter Equal to or Less than Ten Microns in Aerodynamic Diameter (PM<sub>10</sub>)”, May 31, 2000. The August 18, 2002, date passed the criteria for a natural exceptional event, and qualifies for treatment through a Natural Events Action Plan (NEAP).

ADEQ submitted a NEAP to EPA on February 19, 2004, and a PM<sub>10</sub> Maintenance Plan on August 18, 2006. All Best Available Control Measures (BACM) have been adopted and implemented by August 2005. BACM for all significant sources of PM<sub>10</sub> contributing to the PM<sub>10</sub> concentrations in Yuma County include enforcement to prevent traffic and trespass on unpaved Irrigation District canal roads, and measures applicable to windblown dust from agricultural practices, disturbed land, uncovered trucks hauling particulate matter, and vacant lots. A public outreach campaign is also being conducted that includes bilingual brochures, a public service announcement, and videos to explain dust control plans for construction site contractors and agricultural practices.

Additional analyses have been prepared by the Air Quality Division to quantify the emission reductions from the implementation of Agricultural Best Management Practices, which began in Yuma August 1, 2005. This work has been accomplished with the help of Yuma farmers, conservation agents, and Arizona Department of Agriculture personnel.

### ***Western Arizona/Sonora Border Air Quality Study***

The purpose of this study is to determine the sources and movement of air pollutants as well as assess their health impacts on residents of far southwestern Arizona and adjacent regions of Mexico. To accomplish this, ADEQ, in partnership with local, state, federal, and tribal governments, have identified six phases to the study: identifying study requirements and collecting meteorological data; siting study for pollutant monitor locations; monitor deployment; data collection; air quality modeling and health risk assessment. The Air Quality Division will carry out a thorough public outreach program during the study. The first phase is well underway. A total of eight meteorological stations have been installed to acquire data on wind, temperature, relative humidity, solar radiation, atmospheric pressure, and lapse rate. Three stations are in Mexico and five in Arizona. The information acquired during this phase will be used with emissions inventory data and exposure potential to determine where air quality monitors should be sited in the next phase of the study.

Continuous and filter-based monitors were deployed in March 2006 at two “supersites” –sites with a full complement of gaseous and particulate monitors – one in San Luis, Rio Colorado, Sonora, and one in the northeastern part of Yuma. The filter-based particulates samplers and the integrated gaseous pollutant samplers have been run on a one in six day schedule. This monitoring is slated to continue into 2007.

In September – October 2004 ADEQ staff conducted an intensive air monitoring study of PM10 at three sites near unpaved roads in San Luis, Rio Colorado. This study was performed, in part, to provide information to Mexican officials that would help them obtain funds to pave the dirt roads. Results of this month-long study with continuous monitors were maximum four-hour and 24-hour PM10 concentrations of 1078 and 190 ug/m3, respectively, with 45% of the averages exceeding the respective guidelines (150 ug/m3 for each averaging period).



**Figure 6 – Map of Western Arizona/Sonora Border Air Quality Study monitoring locations.**

### **Impacts of Cement Plants on the Sycamore Canyon Wilderness**

One of 12 Class I Wilderness Areas in Arizona – all designated by the Clean Air Act for special protection of their pristine air quality – Sycamore Canyon begins high on the Mogollon Rim 20 miles southwest of Flagstaff and develops into a picturesque, steep-walled canyon that meets the Verde River upstream of Clarkdale. Six miles south of the canyon’s mouth is one of the state’s two large Portland cement plants. About 20 miles west of the canyon is the site of a proposed cement plant. As part of the permit application for the proposed plant, visibility and deposition modeling were performed to assess the impacts of its emissions on Sycamore Canyon. To give the Federal Land Managers a more complete picture, ADEQ staff and a contractor began air quality modeling of the combined (and separate) effects of the two plants on the canyon. This work, which finished in autumn 2005, provided the regulatory agencies with a comprehensive assessment of the present visibility degradation in the canyon from the existing plant, as well as the combined degradation from the two plants.

## ***Joint Air Toxics Assessment Project (JATAP)***

The second phase of the Joint Air Toxics Assessment Project (JATAP) began in January 2005 with a full-year air toxics monitoring program in greater Phoenix. Funding was provided through EPA Region 9 and EPA's Office of Air Quality, Planning and Standards (OAQPS). Carried out through the Institute for Tribal Environmental Professionals (ITEP) in Flagstaff, the JATAP coalition consists of staff from the following agencies and tribes:

- EPA - Region 9
- EPA - Office of Air Quality Planning and Standards
- Salt River Pima - Maricopa Indian Community
- Ft. McDowell Indian Community
- Gila River Indian Community
- Maricopa County Environmental Services Department (MCESD)
- Arizona Department of Environmental Quality (ADEQ)
- Maricopa Association of Governments (MAG)
- Pinal County Air Quality Control District (PCAQCD).

Following a pilot-scale monitoring study to determine which Hazardous Air Pollutants (HAPs) are of most concern in South Phoenix and the Gila River Indian Community in 2003 -2004, this larger air toxics monitoring effort had nine sites: one each in the Gila River, Salt River, and Fort McDowell Indian Communities and five in central and west-central Phoenix. These air toxics concentrations will provide enough information for a preliminary risk assessment, to be conducted in late 2006, and, if funding can be obtained, for a full-scale risk assessment based on an emissions inventory, air quality modeling, and risk assessment modeling. This full-scale assessment is planned for 2007 – 2009.

## ***Regional Haze***

Regional haze is caused by the emissions of air pollutants from a wide variety of sources located over a large geographic area. The haze obscures scenic vistas, which degrades our parks and wilderness areas and interferes with people's enjoyment and recreation in those areas. In 1977, the federal Clean Air Act set a goal to remedy any existing visibility impairment, and prevent any future impairment, from manmade pollution at 158 national parks and wilderness areas known as mandatory Federal Class I areas. The Regional Haze State Implementation Plan (SIP) submitted to EPA in December 2003, focused on four of the 12 national parks and wilderness areas in Arizona: Grand Canyon National Park, Petrified Forest National Park, Sycamore Canyon Wilderness, and Mount Baldy Wilderness. The remaining eight Class I areas will be addressed in a SIP to be submitted to EPA by the December 17, 2007, deadline.

The 2003 Regional Haze SIP relied on a demonstration of how the state is implementing the recommendations of the Grand Canyon Visibility Transport Commission to satisfy reasonable progress toward the national visibility goal. All SIPs from this point on will need to assess the current conditions at a Class I area and then determine what strategies would be necessary

should the area be found to have impaired visibility. Areas with good visibility will need to determine strategies to assure those areas maintain good air quality. Western states developing SIPs under sections 309(g) and 308 of the Federal Regional Haze Rule will have assistance with the assessment and strategies portion of the SIP from the Western Regional Air Partnership (WRAP at [www.wrapair.org](http://www.wrapair.org)).

The Air Quality Division (AQD) has an expanded role regarding regional haze. Extensive fire regulations and policy were developed for the 2003 Regional Haze SIP and the Enhanced Smoke Management Plan will continue to be an important part of regional haze. AQD will perform emissions tracking and modeling necessary to determine specific conditions at Arizona Class I areas beyond what WRAP will provide. Arizona will also implement SO<sub>2</sub> Milestones and Backstop Trading Program, which is a voluntary program for stationary sources emitting 100 tons or more per year of sulfur dioxide that will be integrated into existing permits, and emissions will be tracked annually. The annual emissions for the stationary sources will be reported to WRAP, and every five years, beginning with 2004, emissions will be compiled into a regional Milestone Report. Should a milestone, representing markers on a decreasing regional emissions cap, be exceeded, the backstop trading program would be activated. The possibility of developing a trading program for NO<sub>x</sub> will also be researched by WRAP and ADEQ. Additional information on regional haze can be found at <http://www.wrapair.org/309/index.html>.

### ***Hazardous Air Emergency Response***

Part of the ADEQ multimedia response team, the Hazardous Air Emergency Response (HAER), formerly known as the Hazardous Air Response Team, is called to emergencies by the Emergency Response Unit (ERU) for those incidents that threaten air quality. HAER's objectives are to monitor air quality for public exposure of air pollutants and to provide meteorological support regarding dispersion. This information is provided to the Arizona Department of Health Services or the County Health Department so that appropriate actions can be taken to protect the public. The Team has a fully equipped van with a variety of grab-sampling and continuous sampling air monitoring equipment, field deployable meteorological monitoring equipment including a mini-SODAR to measure winds up to 1,000 feet above the ground, a portable gas chromatograph / mass spectrometer, remotely telemetered video surveillance systems, and many portable particulate monitors for monitoring emissions from fires and other particulates sources. It is staffed by two volunteer members of the Air Quality Division.

Since it started in 1992, HAER (and formerly HART) has responded to 124 incidents. During calendar year 2005, HART responded to five incidents: four wildfires and one dump fire. The wildfires were "BART" near Carefree, "Cave Creek Complex Fire", "Florida" near Tucson, and the "Edge Complex" near Pumpkin Center. The dump fire was at Naco. Through August 2006, HART responded to six incidents: three wildfires ("February" near Payson, "Brins" near Sedona, and "Warm" near Fredonia); the Goodrich UPCO Process Fire, the American Recycling Fire, and the Ecology Recycling Fire.


## Trends

---

### Introduction

Whether air quality meets the standards is a central question – explored at length in the second chapter of this report, but one posed more often is whether it is improving or deteriorating. In Arizona, because of the phasing out of leaded gasoline in the mid-1970s and the installation of effective controls on copper smelters in the 1980s, the concentrations of both lead and SO<sub>2</sub> decreased rapidly.

Although improvements have also been made in the concentrations of CO, O<sub>3</sub> and particulates, O<sub>3</sub> concentrations in the greater Phoenix area are virtually equal to the standard and PM<sub>10</sub> concentrations exceed the standards in Nogales, Phoenix, Buckeye, and Casa Grande. Visibility -- the aspect of the atmosphere most obvious to the population -- has been measured continuously in urban and pristine parts of the state long enough to establish trends. The following discussions examine the trends in these three common air pollutants and visibility in Arizona.




**Figure 7 - Average Best & Average Worst Visibility Impairment in the Phoenix Area**


### Carbon Monoxide

Since the mid to late 1970s, CO concentrations have declined dramatically. In Tucson, the maximum annual eight-hour concentration at 22nd Street and Alvernon declined from 12.0 in 1978 to 2.2 parts per million (ppm) in 2005 – a decrease of 82% (Figure 8).

In Phoenix at 18th Street and Roosevelt (Central Phoenix), the decline was from 23.0 ppm in 1975 to 4.1 ppm in 2005 – a decrease of 82% (Figure 9). The number of exceedances of the eight-hour standard in Phoenix decreased from 75 to 0 at Central Phoenix. The entire Phoenix network of CO monitors recorded over 100 exceedances each year from 1981 through 1986, with an average of 134 per year. The last recorded exceedance was in 1999. Most of this improvement can be attributed to Federal new-vehicle emission standards, augmented by emission reductions from the vehicle inspection and maintenance program, which began in 1976, and the use of oxygenated fuels in the winter, beginning in 1989.



**Figure 8 - Eight-hour carbon monoxide maxima at 22<sup>nd</sup> Street and Alvernon Way in Tucson**



**Figure 9 - Eight-hour carbon monoxide maxima at 18<sup>th</sup> Street and Roosevelt in Central Phoenix**

## Ozone

### One-Hour Ozone Concentrations

Maximum one-hour O<sub>3</sub> concentrations have remained steady in Yuma, but have declined in Phoenix and Tucson since 1980 (Figure 10). These decreases have been 32% and 23%, for Phoenix and Tucson, respectively. The Phoenix decrease in O<sub>3</sub> concentrations since 1980 has been nowhere near as pronounced as its declining CO trend, but the net result has been similar: only one exceedance of the O<sub>3</sub> standard has been recorded after 1996. The one-hour standard was officially declared attained on May 16, 2001. Changes in emissions would not be expected to produce proportional changes in concentration because of the relatively high background level of O<sub>3</sub> and its photochemical formation from hydrocarbons and nitrogen oxides. Yuma and Tucson have met the one-hour standard consistently since monitoring began. In the Phoenix airshed, the standard was exceeded regularly through the mid 1990s, with a gradual decrease to 1996, after which the concentrations remained steady and just below the standard until 2005. In 2005, the network maximum one-hour O<sub>3</sub> concentration increased in the Phoenix area to the exceedance level (but did not constitute a violation, see Chapter 2, 1-hr O<sub>3</sub> standard) at one site in the Phoenix area.




Figure 10 - Maximum one-hour ozone concentrations in three cities

### Eight-Hour Ozone Concentrations

The eight-hour O<sub>3</sub> standard, proposed by EPA in 1997 and officially promulgated in 2003, is expressed as the three-year average of the annual fourth-highest concentration, not to exceed 0.08 parts per million. However, due to instrument precision and rounding, this standard translates into a numerical value of 0.085 ppm: any value 0.085 ppm and above is an exceedance. Long-term trends of the fourth-highest ozone concentrations in Tucson fluctuate between 0.060 and 0.080 ppm, but, overall, are steady (Figure 11).

A similar pattern in eight-hour ozone trends also characterizes Yuma, where, although the values are slightly higher than Tucson's, the nearly constant trend is apparent (Figure 12).




Figure 11 - Annual fourth-highest eight-hour ozone concentrations in Tucson




Figure 12 - Annual fourth-highest eight-hour ozone concentrations in Yuma

In contrast to the within-standard concentrations in Tucson and Yuma, 20 of the 33 sites in greater Phoenix recorded annual fourth-highest O<sub>3</sub> values in excess of 0.084 ppm from 1996 to 2005. In metropolitan Phoenix, there were three sites with an

annual fourth-highest eight-hour O<sub>3</sub> concentration in excess of 0.084 ppm in 2005. On the bright side, elevated values of the annual fourth-highest eight-hour O<sub>3</sub> concentration occurred at fewer monitoring sites and at lower values in 2005 than in 1996, with most of the improvement taking place since the 2000 to 2002 period. For instance, of the nine sites operational both in 1996 and 2005, six recorded fourth-highest values greater than 0.084 ppm in 1996, but there were none in 2005. The values have decreased through time as well, with typical fourth-highest concentrations decreasing from 1996 to 2005: Phoenix Supersite, 0.087 ppm to 0.076 ppm; South Phoenix, 0.084 ppm to 0.076 ppm; South Scottsdale, 0.089 ppm to 0.077 ppm, and North Phoenix, 0.092 ppm to 0.084 ppm.

Looking at the specific statistical form of the standard -- the three-year average of the annual fourth-highest eight-hour ozone concentration -- metropolitan Phoenix did not exceed the standard in the three year period from 2003 to 2005, and, as with the annual fourth-highest values, the extent and severity have been decreasing with time.


ADEQ reviewed the three-year periods ending with 1998 through 2005: the first being 1996 to 1998 and the last being 2003 to 2005. In the first two three-year periods 1996 to 1998, and 1997 to 1999 (Table 23); 13 and five monitoring sites, respectively, had average fourth-highest values equal to or exceeding 0.085 ppm. In the last two periods, the numbers of such sites had decreased to one and zero, respectively. The magnitude of these three-year averages has decreased substantially, as well. The highest average for the period ending in 1998 was 0.0923 ppm; the highest average in 2005 was nine percent lower at 0.084 ppm. These trends are consistent with the decreasing one-hour maximum ozone trends; however, most of the decrease in eight-hour ozone concentrations occurred since 2000, five years later than the decrease in the one-hour concentrations. This suggests that, assuming favorable meteorological conditions, the eight-hour standard may be achieved in two to three years.

**Table 23. Three-Year Averages of the Annual Fourth-Highest Eight-Hour Ozone Concentrations in Phoenix and Environs**

(Units are in parts per million (ppm). Bold values in yellow cells equal or exceed the operational standard of 0.085 ppm)

Site	1996-1998	1997-1999	1998-2000	1999-2001	2000-2002	2001-2003	2002-2004	2003-2005
Emergency Mgmt	<b>0.0873</b>	0.0847	0.0823	0.0763	Closed	Closed	Closed	Closed
North Phoenix	<b>0.0923</b>	<b>0.0880</b>	<b>0.0863</b>	<b>0.0853</b>	<b>0.0857</b>	<b>0.0856</b>	0.0837	0.0830
Salt River Pima	<b>0.0907</b>	0.0843	Closed	Closed	Closed	Closed	Closed	Closed
Phoenix Supersite	<b>0.0853</b>	0.0737	0.0727	0.0723	0.0770	0.0766	0.0743	0.0740
Blue Point	<b>0.0893</b>	<b>0.0860</b>	<b>0.0887</b>	<b>0.0853</b>	0.0843	0.0840	0.0823	0.0800
Apache Junction	<b>0.0860</b>	0.0817	0.0813	0.0797	0.0797	0.0763	0.0737	0.0690
Mesa	<b>0.0853</b>	0.0810	0.0793	0.0773	0.0737	Closed	Closed	Closed
Pinnacle Peak	<b>0.0867</b>	0.0810	0.0817	0.0820	<b>0.0850</b>	0.0840	0.0783	0.0780
Fountain Hills	<b>0.0850</b>	0.0823	0.0817	0.0810	0.0847	0.0840	0.0813	0.0820
Falcon Field	<b>0.0850</b>	0.0823	0.0817	0.0810	0.0800	0.0813	0.0777	0.0750
Mount Ord	<b>0.0907</b>	<b>0.0873</b>	<b>0.0887</b>	0.0847	Closed	Closed	Closed	Closed
South Scottsdale	0.0807	0.0753	0.0760	0.0760	0.0787	0.0783	0.0763	0.0760
West Phoenix	0.0847	<b>0.0853</b>	<b>0.0860</b>	0.0823	0.0800	0.0786	0.0777	0.0720
Maryvale	0.0837	0.0813	0.0830	0.0783	0.0790	0.0800	0.0835	0.0830
Humboldt Mt.	<b>0.0880</b>	<b>0.0860</b>	<b>0.0863</b>	0.0847	<b>0.0850</b>	<b>0.0873</b>	<b>0.0850</b>	0.0840
Tonto Monument					<b>0.0870</b>	<b>0.0855</b>	0.0827	0.0810
Queen Valley				0.0790	0.0810	0.0830	0.0810	0.0810
Cave Creek				0.0830	0.0845	0.0840	0.0817	0.0800
Hillside	<b>0.0855</b>	0.0810	0.0833	0.0810	0.0827	0.0773	0.0777	0.0720
Rio Verde	0.0840	0.0833	0.0837	<b>0.0850</b>	0.0847	0.0837	0.0840	0.0810
West Chandler	0.0820	0.0733	0.0733	0.0747	0.0793	0.0797	0.0770	0.0740
Maximum	<b>0.0923</b>	<b>0.0880</b>	<b>0.0887</b>	<b>0.0853</b>	<b>0.0857</b>	<b>0.0873</b>	<b>0.0850</b>	0.0840
n ≥ 0.085 ppm	13	5	5	3	4	3	1	0

Illustrated in Figure 13 are the three-year averages from nine monitoring sites, which are listed in Table 23, that have a long-term period of operation and have recorded one or more averages above the standard. Although there is considerable site-to-site variability, the overall impression is a distinctly downward trend, perhaps best exemplified by Apache Junction. Humboldt Mountain goes against this trend, but has a downward trend since 2001 to 2003.



**Figure 13 - Phoenix area eight-hour ozone trends: three-year averages of the annual fourth-highest concentrations**

Reviewing these sites together (Figure 14), the maximum value fluctuates at or just above the standard for all of the periods except the first, with a range from 0.085 to 0.088 ppm. The average of these sites, after a steady trend for the first half of the record, moves decidedly down in the latter half. These sites seem capable of producing maximum values at or slightly above the standard throughout the period of record; but their average is displaying a robust decline since 2000 to 2002.

**Figure 14 - Phoenix area eight-hour ozone trends: three-year averages of the annual fourth-highest concentrations, expressed as the average and maximum of nine long-term sites**




If the trend in the latter half of the period continues, attainment of the eight-hour ozone standard may not be far away.


## Particulates

### PM<sub>10</sub>

PM<sub>10</sub> concentrations have decreased considerably throughout the state in both urban and rural settings. Nonetheless, this pollutant, more than any other, continues to exceed its standards. For example, annual PM<sub>10</sub> concentrations in South Phoenix averaged 68.7  $\mu\text{g}/\text{m}^3$  from 1985 to 1987, but only 51.0  $\mu\text{g}/\text{m}^3$  in 2003 to 2005, a decrease of 26 percent, but still over the standard. Similar percentage decreases occurred since the 1980s at Central Phoenix and West Phoenix (Figures 15a & b). Figures 15a & b, which show the three-year moving averages, have two distinct similarities: first, one or more sites shows dramatic improvement in the earliest part of the record; and, second, all sites show improvement in the latter part. In Figure 15a the exceptional site is Chandler which peaks midway through the period of record, but has the same downward trend at the end of the record. Figure 15b shows that South Phoenix increased steadily from the period ending in 1997 through the one ending in 2002, while West Phoenix had only a modest increase during this interval.



**Figure 15a - Three-Year Moving Averages of Annual Average PM<sub>10</sub> at four metropolitan Phoenix sites with moderate PM<sub>10</sub> levels (each data point is the average of the three years ending in that year (e.g. “2005” is the average of 2003, 2004, and 2005)).**



**Figure 15b - Three-Year Moving Averages of PM<sub>10</sub> at four metropolitan Phoenix sites with higher PM<sub>10</sub> concentrations**

Despite these improvements, unlike the trends for CO and O<sub>3</sub>, PM<sub>10</sub> standards continue to be violated. Annual concentrations for the last 14 years, presented in Table 24, demonstrate that some sites in metropolitan Phoenix have been above the standard for one or more years: Chandler, South Phoenix, West Phoenix and Greenwood. Of these four sites there have been 17 exceedances of the annual standard over the last 9 years (1997 to 2005). Each of these sites presents a different mix of localized emission sources. Chandler's emissions have gone from agricultural to earthmoving for residential and road construction. South Phoenix and Durango, near the industrial Salt River area, are influenced by emissions from the industrial sources there. Without any nearby industrial or earthmoving activity, West Phoenix PM<sub>10</sub> concentrations would appear to be the result of the transport of metropolitan wide emissions into this part of town. Two miles southeast of West Phoenix, Greenwood combines the high regional concentrations with its close proximity to a major arterial street and freeway.


**Table 24: Annual PM<sub>10</sub> Concentrations in Metropolitan Phoenix (µg/m<sup>3</sup>)**

<i>Year</i>	<i>Central Phoenix</i>	<i>Chandler</i>	<i>Glendale</i>	<i>North Phoenix</i>	<i>South Phoenix</i>	<i>West Phoenix</i>	<i>Mesa</i>	<i>South Scottsdale</i>	<i>Greenwood</i>	<i>Durango</i>
1992	42	<b>56</b>	34	35	48	47	29	34		
1993	43	<b>58</b>	35	34	44	44	35	34		
1994	43	50	33	35	44	43	36	38		
1995	44	<b>56</b>	33	36	46	44	35	36		
1996	41	<b>62</b>	34	37	47	45	33	35		
1997	44	<b>61</b>	38	38	<b>55</b>	<b>51</b>	43	41	<b>61</b>	
1998	38*	45	29	29	31*	39	29	34	50	
1999	44	<b>60</b>	36	35	49	<b>51</b>	35	40	<b>56</b>	<b>69</b>
2000	46	<b>57</b>	41	37	<b>61</b>	<b>53</b>	37	40	<b>61</b>	<b>70</b>
2001	38	48	33	30	50	43	30	33	49	<b>59</b>
2002	43	<b>56</b>	40	37	<b>60</b>	<b>53</b>	36	37	<b>55</b>	<b>70</b>
2003	40	50	36	34	<b>52</b>	46	34	36	<b>51</b>	<b>62</b>
2004	37	40	26	25	46	37	23	26	44	<b>52</b>
2005	39	49	29	30	<b>55</b>	45	30	34	<b>52</b>	<b>66</b>

**Bold values in yellow cells exceed the annual standard of 50 µg/m<sup>3</sup>.**


**\* Does not satisfy EPA summary criteria of 75% data recovery.**

The highest PM<sub>10</sub> concentrations in metropolitan Phoenix are in southwest Phoenix, along the Salt River from about 7<sup>th</sup> Street to 59<sup>th</sup> Avenue. Although most of the area is industrial, there are many residential areas. The PM<sub>10</sub> record in this area since 1994 is shown in Figure 16. The West 43<sup>rd</sup> Avenue site is the replacement for the Salt River site. Concentrations have exceeded the standard every year of monitoring in this area.



**Figure 16 - Annual PM<sub>10</sub> concentrations in the Salt River area**

In Tucson, the background site of Corona de Tucson and the rural site of Green Valley have had fairly steady trends of PM<sub>10</sub>, but the four long-term urban sites all show substantial decreases since the mid 1980s. Orange Grove had a three year average of 43.3 µg/m<sup>3</sup> in 1985 to 1987, but has since decreased 35 percent to a concentration of 28.3 µg/m<sup>3</sup>. South Tucson, Prince Road and Broadway/Swan showed smaller, but substantial, decreases (Figure 17), with similar patterns of an early decrease, followed by a period of gradual increases, and ending with decreasing trends in the last five years.



**Figure 17 - Three-year moving averages of annual average PM<sub>10</sub> at six metropolitan Tucson sites**

These PM<sub>10</sub> reductions in the urban settings can be attributed to a reduction of coarse particulate emissions from paving roads, alleys and road shoulders, and better controls of dust emissions from construction sites.

Throughout the rest of the state, PM<sub>10</sub> concentrations have declined since 1985 at many sites. Figure 18 presents these trends as three-year moving averages. Consider the trends over the last 20 years for a group of high concentration sites outside of the Phoenix area: Payson and Paul Spur have been reduced by more than 70 percent, Douglas concentrations have been reduced by nearly half, Rillito has decreased 40 percent and Yuma has decreased 26 percent. For most of the sites, nearly all of the improvement took place from the mid 1980s to the mid 1990s. The percentage improvement during this ten-year period varied from 24 to 65%, depending on the site, a remarkable decrease. After this point, two sites continued to decrease (Paul Spur and Payson); three sites (Nogales, Yuma, and Rillito) increased until the early part of 2000's; and two sites have remained about the same (Douglas and Hayden). Between 2001 and 2005 Nogales and Yuma have had a decreasing trend, and Rillito has leveled out. At the beginning of the period, six of the seven sites were above the standard; all have been within the standard since the mid 1990s. In each of these localities, road paving, better industrial dust controls, and (in Payson only) cleaner fireplaces and woodstoves can be given credit for the improvement. All of these PM<sub>10</sub> emission reductions were accomplished through State Implementation Plan activities led by the Air Quality Division.




Figure 18 - Three-year moving averages of the annual average PM<sub>10</sub> concentrations at sites with higher concentrations from different location in the state

PM<sub>10</sub> concentrations at sites with lower concentrations have also decreased with Ajo concentrations reduced by 44 percent, Bullhead City by 56 percent, and Safford by 50 percent. Other sites with lower concentrations at lower elevations were steady or slightly decreasing (Figure 19).





Figure 19 - Three-year moving averages of annual average PM<sub>10</sub> concentrations at lower concentration sites at lower elevations

Low-concentration sites at higher elevations – all within the  $50 \mu\text{g}/\text{m}^3$  annual standard for their periods of record, have also noticeably declined since the mid 1980s. Clarkdale decreased 20 percent; Flagstaff, 50 percent; Prescott, 17 percent; and Show Low, 38 percent. (The site in Prescott was moved to Prescott Valley in 2002.) Part of these decreases can be attributed to cleaner-burning wood stoves and fireplaces (Figure 20). What is encouraging when examining these various sites is that not a single one, whether urban, industrial, agricultural or rural, shows a consistent long-term, upward trend.



**Figure 20 - Three-year moving averages of annual average  $\text{PM}_{10}$  concentrations at sites with low concentration at higher elevations.**

### $\text{PM}_{2.5}$

As was discussed earlier,  $\text{PM}_{2.5}$  has not been monitored as long as  $\text{PM}_{10}$ . Measurements of this fine particle fraction were taken with dichotomous samplers at all sites until the late 1990s, when monitoring with  $\text{PM}_{2.5}$  reference instruments began. The dichotomous samplers give an approximate cutpoint between fine and coarse particles somewhere in the range of 2.5 to 3.0 microns. Consequently, measurements taken with these samplers should be termed “fine particulates” or “ $\text{PM}_{\text{fine}}$ ”, and not “ $\text{PM}_{2.5}$ .” In Arizona, the earliest measurements began in 1991 in the rural cities and towns, 1994 in Tucson, and the following year in Phoenix; these data are presented in Tables 25a, b, and c, and Figures 21, 22, and 23.

Figure 21 shows the three-year moving averages of five sites located at various locations in the state. Douglas and Flagstaff have shown flat trends, while Payson's trend is significantly down by 54 percent. Nogales shows an increase in PM<sub>2.5</sub> concentration from 2004 to 2005, but is still within the standard. Exceedances of the annual PM<sub>2.5</sub> standard occurred for four years in Payson and for one year in Higley (Figure 22). Payson, Nogales, and the central area of Phoenix have the highest concentrations of fine particulates. Flagstaff and the urban fringe of Tucson (the Tangerine and Fairgrounds sites) have the lowest concentrations. Fine particulate trends in metropolitan Phoenix decrease from 1995 through 1998 but increase slightly thereafter through 2005, as seen in Figure 22. Inconsistent with this latter trend, Apache Junction concentrations have decreased steadily since 1999. In metropolitan Tucson (Figure 23), records show that the PM<sub>2.5</sub> concentrations at Orange Grove and Children's Park have decreased significantly since monitoring began and that the Central site increased from 2001 to 2004.

**Table 25a. Annual  $PM_{fine}$  and  $PM_{2.5}$  Concentrations Throughout Arizona (in  $\mu\text{g}/\text{m}^3$ )**

Statewide					
Year	Yuma	Flagstaff	Payson	Nogales	Douglas
1991	7.6	N/A	<b>17.9</b>	12.3	8.5
1992	5.7	N/A	<b>17.2</b>	12.6	7.9
1993	6.1	5.4	13.0	9.7	7.9
1994	8.3	4.9	<b>15.8</b>	10.4	8.1
1995	7.2	5.8	<b>15.7</b>	14.3	7.7
1996	8.7	11.2	14.4	13.3	8.3
1997	6.0	5.0	12.2	11.3	6.0
1998	8.3	4.7	10.9	12.5	6.8
1999	7.9	8.4 *	9.8 *	12.5 *	7.9*
2000	8.7	6.9 *	10.0 *	12.8 *	7.1*
2001	10.0	7.1 *	8.8 *	10.7 *	7.2*
2002	N/A	7.1 *	10.0 *	12.1 *	7.4*
2003	N/A	5.6 *	8.9 *	11.3 *	6.4*
2004	N/A	6.8*	9.5*	10.8*	7.1*
2005	N/A	6.0*	8.3*	13.1*	7.3*

**Bold values in yellow exceed the annual standard of  $15 \mu\text{g}/\text{m}^3$ .**

**N/A – Data are not available.**

**\* Data are from federal reference monitors. not dichot monitors.**

**Table 25b. Annual  $PM_{fine}$  and  $PM_{2.5}$  Concentrations in the Phoenix Metropolitan Area ( $\mu\text{g}/\text{m}^3$ )**

Year	Higley	Tempe	Supersite	ASU West	Estrella	West PHX	Apache Junction
1995	<b>15.4</b>	10.0	12.6	11.1	11.7	N/A	N/A
1996	11.1	10.0	13.4	10.5	11.1	N/A	N/A
1997	10.4	9.8	12.1	9.1	7.9	N/A	N/A
1998	9.4	9.4	10.9	8.3	7.1	N/A	N/A
1999	11.1	10.7 *	12.2 *	9.1	8.9	N/A	7.4 *
2000	10.0	10.3 *	11.4 *	8.5	7.7	13.8 *	7.2 *
2001	N/A	9.3 *	9.2 *	N/A	7.4	10.8 *	6.2 *
2002	N/A	10.3 *	11.6 *	N/A	6.7	12.5*	6.3 *
2003	N/A	9.6 *	11.2 *	N/A	7.3	10.6 *	6.3 *
2004	N/A	N/A	9.7*	N/A	N/A	11.6*	5.5*
2005	N/A	N/A	9.7*	N/A	N/A	12.9*	5.5*

**Bold values in yellow exceed the annual standard of  $15 \mu\text{g}/\text{m}^3$ .**

**N/A – Data are not available.**

**\* Data are from federal reference monitors. not dichot monitors.**

**Table 25c. Annual  $PM_{fine}$  and  $PM_{2.5}$  Concentrations in the Tucson Metropolitan Area ( $\mu\text{g}/\text{m}^3$ )**

Year	Orange Grove	22/Craycroft	Tangerine	Fairgrounds	Central	Children's Park
1994	9.4	7.9	5.3	5.8	8.9	N/A
1995	8.9	8.6	5.3	5.1	8.9	N/A
1996	8.2	6.4	4.9	4.7	7.7	N/A
1997	8.7	7.3	5.1	5.5	8.4	N/A
1998	7.3	6.3	5.0	5.0	7.5	N/A
1999	9.6 *	7.5	N/A	N/A	7.2	8.7 *
2000	7.7 *	N/A	N/A	N/A	7.8	6.8 *
2001	7.6 *	6.0	N/A	N/A	7.6	6.8*
2002	6.3*	8.6	N/A	N/A	8.3	6.6*
2003	6.4*	7.5	N/A	N/A	9.7	6.5*
2004	5.8*	N/A	N/A	N/A	N/A	6.6*
2005	6.3*	N/A	N/A	N/A	N/A	5.9*

**Bold values in yellow exceed the annual standard of  $15 \mu\text{g}/\text{m}^3$ .**

**N/A - Data are not available.**

**\* Data are from federal reference monitors. not dichot monitors.**




Figure 21 - Statewide three-year moving averages of annual averages of PM<sub>2.5</sub>




Figure 22 – Metropolitan Phoenix annual averages of PM<sub>2.5</sub>




Figure 23 – Metropolitan Tucson annual averages of PM<sub>2.5</sub>


### Visibility

Optical measurements of visibility have been made continuously since 1993 in Tucson and since 1994 in Phoenix. Light extinction, the degree to which light is reduced by its interaction with particles and gases in the atmosphere, is measured continuously with transmissometers. These measurements have been divided into six categories: the mean of the dirtiest 20 percent of all hours, the mean of all hours and the mean of the cleanest 20 percent of all hours, for both the entire day and the 5 to 11 a.m. period. The units of measurement are inverse megameters ( $Mm^{-1}$ ): the higher the light extinction value in  $Mm^{-1}$ , the more visibility is reduced. Tables 26a and b, and Figures 24 and 25 present these data.

<b>Table 26a: Annual Average Light Extinction in Phoenix (<math>Mm^{-1}</math>)</b>						
<b>Year</b>	<b>All Hours</b>			<b>5-11 a.m.</b>		
	<b>Dirtiest 20%</b>	<b>Mean</b>	<b>Cleanest 20%</b>	<b>Dirtiest 20%</b>	<b>Mean</b>	<b>Cleanest 20%</b>
1994	N/A	64	29	N/A	70	33
1995	141	77	38	137	80	43
1996	134	78	43	130	80	45
1997	131	81	48	136	87	53
1998	133	78	45	136	84	50
1999	127	72	38	128	77	42
2000	131	74	38	134	80	42
2001	118	69	36	118	73	42
2002	124	75	42	125	79	46
2003	131	72	36	135	78	42
2004	121	69	35	126	75	42
2005	126	72	36	128	78	43

<b>Table 26b: Annual Average Light Extinction in Tucson (<math>Mm^{-1}</math>)</b>						
<b>Year</b>	<b>All Hours</b>			<b>5-11 a.m.</b>		
	<b>Dirtiest 20%</b>	<b>Mean</b>	<b>Cleanest 20%</b>	<b>Dirtiest 20%</b>	<b>Mean</b>	<b>Cleanest 20%</b>
1993	101	60	34	139	74	37
1994	95	59	36	109	68	41
1995	104	62	35	116	69	38
1996	99	62	37	113	71	40
1997	93	60	36	108	68	38
1998	102	57	28	119	69	34
1999	90	57	35	107	65	38
2000	98	56	27	114	66	31
2001	96	55	26	109	66	33
2002	87	49	24	109	61	29
2003	88	52	26	107	62	30
2004	97	58	27	113	67	32
2005	101	61	31	125	76	39


Distinct trends from these tabular data are somewhat difficult to discern, partly because of the year-to-year variability and partly because the long-term changes for most categories are rather small. Rather than plotting all of these data, this report is limited to the “all hours” categories, since both the “5-11 a.m.” and “all hours” trends are virtually identical. In Figures 24 and 25 these light extinction data have been plotted as three-year moving averages. The first year shown, 1996, is the average of 1994, 1995, and 1996, and so on.



**Figure 24 – Light extinction trends for Phoenix, shown as three-year moving averages, for all hours**

Considering Phoenix first, the steady improvement through 2002 in the 20% dirtiest category is evident. The most recent period (2003 to 2005) in this category is 7% lower than the first full three-year period. For both the mean and 20% cleanest days, however, the steadily downward trend of the dirtiest 20% category is replaced by a more complicated trend – one in which the first two three-year periods increase through 1998, but the subsequent periods gradually decrease and eventually level out by 2001. What’s happened in this twelve-year period (1994 to 1996 compared to 2003 to 2005) is that visibility has gotten somewhat better with a 7% decrease for the dirtiest 20%, and a 3% decrease for both the mean and cleanest 20%. There is, however, a slight rise in the early years for the mean and 20% cleanest and another very gradual increase over the last three periods for the dirtiest 20% (as seen in the figure above and in Table 26a).


Unlike Phoenix, visibility in Tucson has improved over the 13-year period when considering the three-year averages for all three statistics: the dirtiest, the mean, and the cleanest (Figure 25). The improvement in the 20% dirtiest days was 5%, which is 2% less than the improvement in Phoenix, but considerably greater improvement has been realized in the 20% cleanest category with a 20% decrease. Somewhat disturbing, however, is the upward trend in the annual statistics from 2002 through 2005 (evident in Table 26b and in the upward trending curves for the last two periods).



**Figure 25 – Light extinction trends for Tucson, shown as three-year moving averages, for all hours**


Since it's impossible for an observer to distinguish between the various grades of the cleanest 20%, perhaps the overall Phoenix-Tucson trends appear the same to their respective residents. That is, over this 12 or 13 year period, there has been a 7% decrease in the light extinction values for the dirtiest days in Phoenix and 5% in Tucson. Residents of each metropolitan area, then, have observed improved visibility for these haziest of days. While the worst of the brown clouds are still quite evident, especially on winter mornings, their frequency and severity over both cities have diminished slightly.

An interesting intercity trend (Figure 26) appears in the cleanest 20% category, where, in the first years of monitoring, Tucson and Phoenix had equal values. As the 1990s progressed, however, Tucson's cleanest days grew decidedly cleaner, while Phoenix's cleanest days had increased light extinction for the first half of the period, followed by a gradual decrease and leveling off in the later part of the record. The result is that in 2003 – 2005, Tucson's cleanest days were 21% cleaner than in Phoenix (28 Mm<sup>-1</sup> vs. 36 Mm<sup>-1</sup>).




**Figure 26 – Light extinction trends for all hours for Phoenix and Tucson, shown as three-year moving averages, for all hours and the cleanest 20% days**

Seasonal patterns also vary between the two cities, with the mean and dirtiest 20 percent of all hourly light extinction values in Phoenix showing more pronounced winter and fall maxima than the Tucson counterparts (Figure 27). Both cities show little seasonal variation in the cleanest 20 percent of all hours. The seasonal light extinction values in Phoenix are considerably higher than Tucson's: for the dirtiest 20 percent of all hours, 52 percent higher in winter, 19 percent higher in spring, 13 percent higher in summer and 49 percent higher in fall. The poorer visibility in Phoenix comes as no surprise to those Arizonans familiar with both airsheds.



**Figure 27 – Seasonal variation in light extinction of the 20% cleanest and 20% dirtiest days in Tucson and Phoenix**

In the following, final, discussion of visibility, light scattering is compared between the urban and rural areas of the state (Figure 28). In each statistical category rural light scattering is considerably lower than the urban. On the dirtiest 20% days, light scattering values in Phoenix are 3.5 times higher than in the rural areas, while values in Tucson are nearly twice as high. Values for the mean and 20% cleanest days show similar results. An interesting comparison between urban and rural areas is that the light scattering values on the worst 20% days in the rural areas are roughly equal to the mean of the urban areas.



**Figure 28 – Comparison of light scattering on the 20% cleanest, mean, and 20% dirtiest days for urban and rural areas.**

## Conclusions

Since monitoring of air pollutants began in the late 1960s in Arizona, considerable progress has been made in reducing concentrations of lead, SO<sub>2</sub>, and CO. Lead has been reduced to near background levels; SO<sub>2</sub> concentrations near copper smelters, which chronically exceeded the standards until the mid-1980s, are now well within these standards; and CO concentrations, which regularly exceeded standards in neighborhoods and near busy intersections in Phoenix (and to a far lesser extent in Tucson), now meet the standards. One-hour O<sub>3</sub> concentrations in Phoenix have met the standard since 1997, the first years since monitoring began. Phoenix one-hour ozone concentrations in the 1980s and early 1990s ranged as high as 0.18 parts per million (the standard is 0.12 ppm), in contrast to the highest, most recent reading of 0.14 ppm in 1996. In 1995-1997, 12 monitoring sites in greater Phoenix exceeded the eight-hour O<sub>3</sub> standard; in 2003 - 2005 no sites exceeded the standard.

Elevated concentrations of PM<sub>10</sub> have been reduced substantially since the mid-1980s, with decreases of 20 to 70 percent in the urban areas and in most rural cities and towns. In Payson and at some industrial sites, PM<sub>10</sub> concentrations have been reduced by as much as two-thirds. By 2005, monitored violations of the PM<sub>10</sub> standard -- a once common occurrence at many sites only ten years ago -- were limited to a few sites in southwest Phoenix, Pinal County, and in Nogales. The severity of the PM<sub>10</sub> problems in these areas, exemplified by the 32 expected exceedances of the 24-hour standard in southwest Phoenix in 2005, point out the need for further controls on emissions. Fine particulates concentrations (PM<sub>2.5</sub>) have decreased in Phoenix and Tucson since the mid 1990s; for example, at the centrally located Phoenix Supersite, the decrease has been 21 percent; at 22nd and Craycroft, in east-central Tucson, the decrease has been 24 percent. Fine particulate trends in rural Arizona, however, have not shown consistency from site to site: Nogales has been steady; Yuma and Flagstaff have increased (by 48 and 26%, respectively); and Douglas and Payson have decreased (by 14 and 48%, respectively).

In spite of the continued growth in Arizona, not a single air pollutant at any site shows a consistent upward trend. Most standards are met all of the time, with the exceptions being the eight-hour O<sub>3</sub> standard on occasional summer days in Phoenix and the PM<sub>10</sub> standards on both an episodic and annual basis at those sites affected by localized dense emissions. This improved air quality -- resulting from emission control programs at the federal, state and local levels -- has benefited the respiratory health of the citizenry and can be considered a consequence of the public support for a cleaner environment.



## Appendix 1 – Site Index

<i>Site Index - Ambient Air Monitoring Locations in Arizona in 2006</i>										
City/Site and Address	Lat.	Long.	Elev. (feet)	Parameters Measured	Operator	Classification	Scale	Objective	AAAD ID Number	AQS ID Number
<b>Apache County</b>										
Greer Water Treatment Plant (Mt Baldy)	34.06	-109.44	8252	Bscat, MET, IMPROVE	ADEQ, USFS	Class I	Regional	Visibility	16323	N/A
Petrified Forest NP	34.82	-109.89	5796	Bscat, MET, IMPROVE, O <sub>3</sub>	NPS	Class I	Regional	Visibility	16473	04-017-0119
TEP – Springerville - Coalyard	34.33	-109.15	6898	PM <sub>10</sub>	TEP	SPM	Unknown	Source Impact	16637	N/A
TEP – Springerville - Coyote Hills	34.17	-109.23	6599	NO <sub>2</sub> , PM <sub>10</sub> , SO <sub>2</sub>	TEP	SPM	Unknown	Source Impact	16638	N/A
<b>Cochise County</b>										
Chiricahua NM Entrance Station (3.5 miles west of monument headquarters)	32.01	-109.39	5130	CASTNET, Bscat, IMPROVE, MET, O <sub>3</sub>	NPS	Class I	Regional	Visibility	16679	04-003-8001
Douglas Red Cross (1445E 15th St.)	31.35	-109.54	4100	IMPROVE, PM <sub>10</sub> , PM <sub>2.5</sub>	ADEQ	SLAMS (PM <sub>10</sub> , PM <sub>2.5</sub> ), Class I	Neighborhood	Population	16503	04-003-1005

<b>Site Index - Ambient Air Monitoring Locations in Arizona in 2006</b>										
<b>City/Site and Address</b>	<b>Lat.</b>	<b>Long.</b>	<b>Elev. (feet)</b>	<b>Parameters Measured</b>	<b>Operator</b>	<b>Classification</b>	<b>Scale</b>	<b>Objective</b>	<b>AAAD ID Number</b>	<b>AQS ID Number</b>
Muleshoe Ranch (Galiuro Wilderness) Closed June 2005	32.33	-110.24	4398	Bscat, MET	ADEQ	Class I	Regional	Visibility	16412	N/A
Naco Border Station (218 1st St.)	31.35	-109.74	4100	Bscat	ADEQ	SPM	Neighborhood	Population	16392	N/A
Paul Spur Chemical Lime Plant	31.37	-109.73	4192	PM <sub>10</sub> , MET	ADEQ	SLAMS (PM <sub>10</sub> )	Middle	Source Impact	16391	04-003-0011
<b>Coconino County</b>										
Flagstaff Middle School (755 N. Bonito)	35.21	-111.65	6904	PM <sub>10</sub> , PM <sub>2.5</sub>	ADEQ	SLAMS	Neighborhood	Population	16707	04-005-1008
Grand Canyon NP Hance (South Rim, 2.5 miles west of village)	35.97	-111.98	7436	O <sub>3</sub> , MET, Bscat, IMPROVE, CASTNET	NPS	Class I	Regional	Visibility	16682	N/A

<b>Site Index - Ambient Air Monitoring Locations in Arizona in 2006</b>										
<b>City/Site and Address</b>	<b>Lat.</b>	<b>Long.</b>	<b>Elev. (feet)</b>	<b>Parameters Measured</b>	<b>Operator</b>	<b>Classification</b>	<b>Scale</b>	<b>Objective</b>	<b>AAAD ID Number</b>	<b>AQS ID Number</b>
Grand Canyon NP Indian Garden (4.5 miles from Bright Angel trailhead)	36.08	-112.13	3795	IMPROVE, Bscat,	NPS	Class I	Regional	Visibility	16683	N/A
Ike's Backbone (Pine Mountain Wilderness)	34° 20'	111° 40'	5232	IMPROVE, Bscat	ADEQ, USFS	Class I	Regional	Visibility	16421	N/A
SRP – Page – Navajo Generating Station (3 miles east of Page)	36.91	-111.39	3647	O <sub>3</sub> , NO <sub>2</sub> , PM <sub>10</sub> , SO <sub>2</sub>	SRP	SPM	Urban	Source Impact	16634	N/A
Sedona Post Office (190 W. Highway 89A)	34.87	-111.77	4195	PM <sub>10</sub>	ADEQ	SPM	Neighborhood	Population	16512	04-005-1010
Sycamore Canyon Camp Raymond	35.14	-111.97	6691	Bscat, IMPROVE, MET	ADEQ, NPS	Class I	Regional	Visibility	16476	N/A
<b>Gila County</b>										
ASARCO - Globe Highway	33.00	-110.77	1948	SO <sub>2</sub>	ASARCO	SPM	Regional	Source Impact	16593	N/A

<i>Site Index - Ambient Air Monitoring Locations in Arizona in 2006</i>										
City/Site and Address	Lat.	Long.	Elev. (feet)	Parameters Measured	Operator	Classification	Scale	Objective	AAAD ID Number	AQS ID Number
ASARCO - Hayden - Garfield AVE	33.00	-110.78	2089	SO <sub>2</sub>	ASARCO	SPM	Neighborhood	Source Impact	16590	N/A
ASARCO - Montgomery Ranch	33.01	-110.80	2326	SO <sub>2</sub>	ASARCO	SPM	Regional	Source Impact	16591	N/A
Hayden - Old Jail (Canyon Drive)	33.01	-110.79	2050	PM <sub>10</sub> , SO <sub>2</sub>	ADEQ, ASARCO	SLAMS (ADEQ SO <sub>2</sub> and PM <sub>10</sub> ) SPM (ASARCO SO <sub>2</sub> )	Neighborhood	Source Impact	16326	04-007-1001
PDMI - Miami - Golf Course	33.41	-110.83	3319	PM <sub>10</sub>	PDMI	SPM	Neighborhood	Source Impact	16629	04-007-8000
PDMI - Miami - Jones Ranch (Cherry Flats Rd.)	33.39	-110.87	4093	SO <sub>2</sub>	PDMI	SPM	Neighborhood	Source Impact	16631	N/A
PDMI - Miami - Town Site (Sullivan St.)	33.40	-110.87	3388	SO <sub>2</sub>	PDMI	SPM	Neighborhood	Source Impact	16632	N/A
Miami - Ridgeline (4030 Linden St.)	33.40	-110.86	3559	PM <sub>10</sub> , SO <sub>2</sub>	ADEQ, PDMI	SLAMS (ADEQ SO <sub>2</sub> ) SPM (PDMI PM <sub>10</sub> )	Neighborhood	Source Impact	16382	04-007-0009

<b>Site Index - Ambient Air Monitoring Locations in Arizona in 2006</b>										
<b>City/Site and Address</b>	<b>Lat.</b>	<b>Long.</b>	<b>Elev. (feet)</b>	<b>Parameters Measured</b>	<b>Operator</b>	<b>Classification</b>	<b>Scale</b>	<b>Objective</b>	<b>AAAD ID Number</b>	<b>AQS ID Number</b>
Payson Well Site (204 W. Aero Dr.)	34.23	-111.33	4910	PM <sub>10</sub> , PM <sub>2.5</sub>	ADEQ	SLAMS	Neighborhood	Population	16317	04-007-0008
Pleasant Valley - Ranger Station (Sierra Ancha USFS Wilderness)	34.09	110.94	5133	IMPROVE, Bscat, MET	ADEQ, USFS	Class I	Regional	Visibility	16446	N/A
Tonto NM (Tonto Natl Forest)	33.65	-111.11	2460	IMPROVE, NO <sub>TL</sub> , O <sub>3</sub>	ADEQ, USFS	Class I	Regional	Visibility	16447	04-007-0010
<b>Graham County</b>										
Safford (523 Tenth Ave.)	32.83	-109.72	2949	PM <sub>10</sub>	ADEQ	SLAMS	Neighborhood	Population	16508	04-009-0001
<b>La Paz County</b>										
Alamo Lake	34.24	-113.56	1282	NO <sub>TL</sub> , O <sub>3</sub>	ADEQ	SLAMS	Regional	Background	34961	04-012-8000
<b>Maricopa County</b>										
ADEQ Building (1110 W Washington)	33.45	-112.09	1082	Visibility (camera)	ADEQ	SPM (Urban Haze)	Urban	Urban Haze	21737	N/A

<i>Site Index - Ambient Air Monitoring Locations in Arizona in 2006</i>										
<b>City/Site and Address</b>	<b>Lat.</b>	<b>Long.</b>	<b>Elev. (feet)</b>	<b>Parameters Measured</b>	<b>Operator</b>	<b>Classification</b>	<b>Scale</b>	<b>Objective</b>	<b>AAAD ID Number</b>	<b>AQS ID Number</b>
Banner Mesa Medical Center (525 W Brown AVE)	33.43	-111.84	1489	Visibility (camera & Bext)	ADEQ	SPM (Urban Haze)	Urban	Urban Haze	19489	N/A
Bethune Elementary School (1310 S. 15th Ave.)	33.44	-112.09	1063	PM <sub>10</sub> , Speciated PM <sub>2.5</sub>	ADEQ	SPM, STN	Neighborhood	Population	17786	04-013-8006
Blue Point (Usery Pass and Bush Highway)	33.55	-111.61	1574	MET, O <sub>3</sub>	MCESD	SLAMS (MET) NAMS (O <sub>3</sub> )	Urban	Maximum Concentration	16417	04-013-9702
Buckeye (SR 85 & Buckeye RD)	33.37	-112.62	840	CO, MET, NO <sub>2</sub> , O <sub>3</sub> , PM <sub>10</sub>	MCESD	SLAMS	Neighborhood	Population	21525	04-013-4011
Cave Creek (37109 N. Lava Lane)	33.83	-112.02	1916	MET, O <sub>3</sub>	MCESD	SLAMS	Urban	Maximum Concentration	16368	04-013-4008
Central Phoenix (1845 E. Roosevelt)	33.46	-112.04	1115	CO, MET, NO <sub>2</sub> , O <sub>3</sub> , PM <sub>10</sub> , SO <sub>2</sub>	MCESD	SLAMS (MET) NAMS (CO, NO <sub>2</sub> , O <sub>3</sub> , PM <sub>10</sub> , SO <sub>2</sub> )	Neighborhood	Population	16329	04-013-3002

<i>Site Index - Ambient Air Monitoring Locations in Arizona in 2006</i>										
<b>City/Site and Address</b>	<b>Lat.</b>	<b>Long.</b>	<b>Elev. (feet)</b>	<b>Parameters Measured</b>	<b>Operator</b>	<b>Classification</b>	<b>Scale</b>	<b>Objective</b>	<b>AAAD ID Number</b>	<b>AQS ID Number</b>
Chandler 1475 E Pecos Rd Closed 12/31/05 Repl. by Higley	33° 17'	111° 49'	1171	MET, PM <sub>10</sub>	MCESD	SLAMS (MET) NAMS (PM <sub>10</sub> )	Neighborhood	Population	16369	04-013-0021
Durango Complex 2702 AC Esterbrook Blvd.	33.43	-112.12	1574	MET, PM <sub>10</sub> , PM <sub>2.5</sub>	MCESD	SLAMS	Middle	Maximum Concentration	16375	04-013-9812
Dysart 16825 N Dysart	33.64	-112.34	1099	CO, O <sub>3</sub> , ADEQ(Bscat)	MCESD ADEQ	SPM, Bscat (Urban Haze)	Neighborhood	Population	19550	04-013-4010
Estrella 15099 W. Casey Abbott Dr., Goodyear	33.38	-112.37	1000	Bscat	ADEQ	SPM (Urban Haze)	Neighborhood	Population	16506	04-013-8005
Estrella Community College 3000 N Dysart Rd.	33.48	-112.35	1000	Visibility (camera)	ADEQ	SPM (Urban Haze)	Urban	Urban Haze	21736	N/A
Falcon Field (4530 E. McKellips)	33.45	-111.73	1017	MET, O <sub>3</sub>	MCESD	SLAMS	Urban	Population	16381	04-013-1010

<i>Site Index - Ambient Air Monitoring Locations in Arizona in 2006</i>										
<b>City/Site and Address</b>	<b>Lat.</b>	<b>Long.</b>	<b>Elev. (feet)</b>	<b>Parameters Measured</b>	<b>Operator</b>	<b>Classification</b>	<b>Scale</b>	<b>Objective</b>	<b>AAAD ID Number</b>	<b>AQS ID Number</b>
Fountain Hills (16426 E. Palisades)	33.61	-111.72	1443	MET, O <sub>3</sub>	MCESD	SLAMS (MET) NAMS (O <sub>3</sub> )	Neighborhood	Maximum Concentration	16376	04-013-9704
Glendale (6000 W. Olive)	33.57	-112.19	1171	CO, MET, O <sub>3</sub> , PM <sub>10</sub>	MCESD	SLAMS (CO, MET, O <sub>3</sub> ), NAMS (PM <sub>10</sub> )	Neighborhood	Population	16378	04-013-2001
Greenwood (I-10 and 27th Avenue)	33.46	-112.12	1109	CO, MET, NO <sub>2</sub> , PM <sub>10</sub>	MCESD	SLAMS	Microscale	Maximum Concentration	16372	04-013-3010
Higley (15500 S. Higley Rd.)	33.31	-111.72	1250	MET, PM <sub>10</sub>	MCESD	SLAMS (MET) SPM (PM <sub>10</sub> )	Neighborhood	Population	16505	04-013-4006
Humboldt Mountain (Pine Mountain Wilderness)	33.98	-111.80	5228	O <sub>3</sub>	MCESD	SLAMS	Regional	Background/ Transport	16416	04-013-9508
JLG Supersite (4530 N. 17 Ave.)	33.50	-112.10	1135	Bscat, CO, NO <sub>2</sub> , Met, O <sub>3</sub> , PM <sub>10</sub> , PM <sub>2.5</sub> , VOC, Speciated PM <sub>2.5</sub>	ADEQ	SPM (Urban Haze) SLAMS (CO, NO <sub>2</sub> , O <sub>3</sub> , PM <sub>2.5</sub> ) PAMS (Type 2) STN	Neighborhood	Population	16328	04-013-9997
Maryvale 6180 W Encanto Closed 03/31/05	33° 28'	112° 20'	1050	CO, O <sub>3</sub> , PM <sub>10</sub>	MCESD	SLAMS	Neighborhood	Population	16379	04-013-3006

<i>Site Index - Ambient Air Monitoring Locations in Arizona in 2006</i>										
City/Site and Address	Lat.	Long.	Elev. (feet)	Parameters Measured	Operator	Classification	Scale	Objective	AAAD ID Number	AQS ID Number
Mesa (370 S. Brooks)	33.41	-111.86	1220	CO, MET, PM <sub>10</sub> , PM <sub>2.5</sub>	MCESD	SLAMS	Neighborhood	Population	16380	04-013-1003
Mesa City Building (Lewis & Main)	33.42	-111.83	1378	Bext	ADEQ	SPM (Urban Haze)	Urban	Urban Haze	19686	N/A
North Mountain Summit (North Mountain)	33.59	-112.07	1640	Visibility (camera)	ADEQ	SPM (Urban Haze)	Urban	Urban Haze	16480	N/A
North Phoenix (601 E. Butler)	33.56	-112.07	1243	CO, MET, O <sub>3</sub> , PM <sub>10</sub>	MCESD	SLAMS	Neighborhood	Population	16390	04-013-1004
Phoenix Transmissometer Receiver (3600 N 2nd AVE)	33.49	-112.08	0	Bext	ADEQ	SPM (Urban Haze)	Urban	Urban Haze	16829	N/A
Phoenix Transmissometer Transmitter (2000 W Bethany RD)	33.53	-112.10	1115	Bext	ADEQ	SPM (Urban Haze)	Urban	Urban Haze	16330	N/A
Pinnacle Peak (25000 N. Windy Walk)	33.71	-111.85	2624	MET, O <sub>3</sub>	MCESD	SLAMS	Urban	Maximum Concentration	16406	04-013-2005

<i>Site Index - Ambient Air Monitoring Locations in Arizona in 2006</i>										
City/Site and Address	Lat.	Long.	Elev. (feet)	Parameters Measured	Operator	Classification	Scale	Objective	AAAD ID Number	AQS ID Number
Rio Verde (25608 N. Forest Rd.)	33.72	-111.67	1640	O <sub>3</sub>	MCESD	SLAMS	Urban	High Downwind Concentration	16396	04-013-9706
South Phoenix (33 W. Tamarisk)	33.4	-112.07	1082	CO, MET, O <sub>3</sub> , PM <sub>10</sub>	MCESD	NAMS (PM <sub>10</sub> ) SLAMS (CO, MET, O <sub>3</sub> )	Neighbor- hood	Population	16377	04-013-4003
South Scottsdale (2857 N. Miller)	33.48	-111.92	1227	CO, MET, NO <sub>2</sub> , O <sub>3</sub> , PM <sub>10</sub> , SO <sub>2</sub>	MCESD	SLAMS (CO, MET) NAMS (NO <sub>2</sub> , O <sub>3</sub> , PM <sub>10</sub> , SO <sub>2</sub> )	Urban/ Neighbor- hood	Population	16398	04-013-3003
Tempe (1525 S College AVE)	33.41	-111.94	1181	CO, MET, NO <sub>2</sub> , O <sub>3</sub>	MCESD	SPM	Neighbor- hood	Population	16405	04-013-4005
Tempe Community Center (3340 S. Rural Rd.) Closed 07/26/2005	33° 23'	111° 55'	1110	PM <sub>2.5</sub>	ADEQ	SLAMS	Neighbor- hood	Population	16509	04-013-9990
Vehicle Emissions Laboratory (600 N 40 <sup>th</sup> ST)	33.46	-1112.0	1050	MET, Bscat	ADEQ	SPM Urban Haze (Bscat)	Urban	Meteorology	16363	04-013-9998

<b>Site Index - Ambient Air Monitoring Locations in Arizona in 2006</b>										
<b>City/Site and Address</b>	<b>Lat.</b>	<b>Long.</b>	<b>Elev. (feet)</b>	<b>Parameters Measured</b>	<b>Operator</b>	<b>Classification</b>	<b>Scale</b>	<b>Objective</b>	<b>AAAD ID Number</b>	<b>AQS ID Number</b>
West Chandler (163 S. Price)	33.3	-111.88	1181	CO, MET, O <sub>3</sub> , PM <sub>10</sub>	MCESD	SLAMS	Neighborhood	Population	16478	04-013-4004
West Forty Third (3940 W Broadway)	33.41	-112.14	1030	MET, PM <sub>10</sub> , Speciated PM <sub>2.5</sub>	MCESD	SPM (PM <sub>10</sub> ), STN	Neighborhood	Maximum Concentration	16659	04-013-4009
West Indian School (3315 W. Indian School Rd.)	33.49	-112.13	1115	CO, MET	MCESD	NAMS (CO) SLAMS (MET)	Micro	Maximum Concentration/ Source Impact	16393	04-013-0016
West Phoenix (3847 W. Earll)	33.48	-112.14	1096	CO, MET, NO <sub>2</sub> , O <sub>3</sub> , PM <sub>10</sub> , PM <sub>2.5</sub> , Speciated PM <sub>2.5</sub>	ADEQ, MCESD	SPM (ADEQ PM <sub>2.5</sub> ) SLAMS (MET, NO <sub>2</sub> , O <sub>3</sub> ) NAMS (CO, PM <sub>10</sub> ), STN	Neighborhood	Population	16477	04-013-0019
<b>Mohave County</b>										
Bullhead City (990 Hwy 95)	35.15	-114.57	561	PM <sub>10</sub>	ADEQ	SLAMS	Neighborhood	Population	16365	04-015-1003
Meadview	35.98	-114.07	2959	Bscat, MET, IMPROVE	USFS	Class I	Regional	Visibility	21298	N/A
Kingman - Praxair NE #1 (I-40 and Griffith Road)	35.03	-114.03	2401	PM <sub>10</sub>	Praxair	SPM	Middle	Source Impact	16554	N/A

<b>Site Index - Ambient Air Monitoring Locations in Arizona in 2006</b>										
<b>City/Site and Address</b>	<b>Lat.</b>	<b>Long.</b>	<b>Elev. (feet)</b>	<b>Parameters Measured</b>	<b>Operator</b>	<b>Classification</b>	<b>Scale</b>	<b>Objective</b>	<b>AAAD ID Number</b>	<b>AQS ID Number</b>
Kingman - Praxair SW #2 (I-40 and Griffith Road)	35.03	-114.14	2358	PM <sub>10</sub>	Praxair	SPM	Middle	Source Impact	16555	N/A
<b>Navajo County</b>										
Petrified Forest NP (1 mile north of park headquarters)	35° 05'	109° 46'	5778	Bscat, IMPROVE, MET, O <sub>3</sub>	NPS	Class I	Regional	Visibility	16473	04-017-0119
Show Low (561 E Deuce of Clubs)	34.25	-110.04	6311	PM <sub>10</sub>	ADEQ	SLAMS	Neighborhood	Population	16603	04-017-0007
<b>Pima County</b>										
22nd St. & Alvernon (3895 E. 22nd)	32.21	-110.91	2516	CO	PDEQ	NAMS	Micro	Maximum Concentration	16676	04-019-1014
22nd St. & Craycroft (1237 S. Beverly)	32.2	-110.88	2581	Bscat, CO, O <sub>3</sub> , NO <sub>2</sub> , SO <sub>2</sub>	ADEQ, PDEQ	SPM (ADEQ Urban Haze Bscat) SLAMS (PDEQ CO, O <sub>3</sub> , NO <sub>2</sub> , SO <sub>2</sub> )	Neighborhood	Population	16410	04-019-1011
Ajo (Well Road)	32.38	-112.86	1801	PM <sub>10</sub> , MET	ADEQ	SLAMS (PM <sub>10</sub> )	Neighborhood	Population	16316	04-019-0001

<i>Site Index - Ambient Air Monitoring Locations in Arizona in 2006</i>										
City/Site and Address	Lat.	Long.	Elev. (feet)	Parameters Measured	Operator	Classification	Scale	Objective	AAAD ID Number	AQS ID Number
Broadway & Swan (4625 E. Broadway)	32.22	-110.89	2516	PM <sub>10</sub>	PDEQ	NAMS	Middle	Maximum Concentration	16550	04-019-1023
Cherry & Glenn (2745 N. Cherry)	32.26	-110.95	2401	CO	PDEQ	SPM	Neighborhood	Population	16675	04-019-1021
Children's Park (400 W. River Rd.)	32.3	-110.98	2286	Bscat, CO, NO <sub>2</sub> , O <sub>3</sub> , PM <sub>2.5</sub> , Speciated PM <sub>2.5</sub>	ADEQ, PDEQ	SPM ( PM <sub>2.5</sub> & ADEQ Urban Haze Bscat) SLAMS ( NO <sub>2</sub> , O <sub>3</sub> ) NAMS (CO), STN	Urban, Neighborhood	Population	16551	04-019-1028
Coachline (9597 N Coachline Blvd)	32.38	-111.13	2227	O <sub>3</sub> , PM <sub>2.5</sub>	PDEQ	SPM	Neighborhood	Population	21580	04-019-1034
Corona De Tucson (22000 S. Houghton Rd.)	32.0	-110.79	3077	PM <sub>10</sub>	PDEQ	SLAMS (PDEQ)	Regional	Background	16677	04-019-0008
Geronimo (2498 N. Geronimo)	32.25	-110.97	2578	PM <sub>10</sub>	PDEQ	SPM (For AQI Purposes Only)	Neighborhood	Population	16678	N/A

<i>Site Index - Ambient Air Monitoring Locations in Arizona in 2006</i>										
<b>City/Site and Address</b>	<b>Lat.</b>	<b>Long.</b>	<b>Elev. (feet)</b>	<b>Parameters Measured</b>	<b>Operator</b>	<b>Classification</b>	<b>Scale</b>	<b>Objective</b>	<b>AAAD ID Number</b>	<b>AQS ID Number</b>
Golf Links & Kolb (2601 S. Kolb Rd)	32.19	-110.84	2660	CO	PDEQ	SPM	Neighborhood	Population	19531	04-019-1031
Green Valley (601 N. La Canada Dr.)	31.88	-110.99	2903	O <sub>3</sub> , PM <sub>2.5</sub> PM <sub>10</sub>	PDEQ	SLAMS	Neighborhood	Population Exposure	16685	04-019-1030
Orange Grove (3401 W. Orange Grove Road)	32.32	-111.04	2175	PM <sub>10</sub> , PM <sub>2.5</sub>	PDEQ	SLAMS (PDEQ PM <sub>10</sub> , PM <sub>2.5</sub> )	Neighborhood	Maximum Concentration/ Population	16510	04-019-0011
Organ Pipe Cactus NM (1 mile SSW of visitor center)	31.95	-112.80	1847	PM <sub>10</sub> , IMPROVE, Bscat	ADEQ	SLAMS (PM <sub>10</sub> )	Regional	Background/ Transport, Visibility	16681	04-019-0005
Prince Road (1016 W. Prince Rd.)	32.27	-110.99	2316	PM <sub>10</sub>	PDEQ	NAMS	Micro	Source Impact	16597	04-019-1009
Rillito (8820 W. Water)	32.42	-111.15	2053	PM <sub>10</sub>	ADEQ, APCC	SLAMS (ADEQ) SPM (APCC)	Neighborhood	Source Impact	16499	04-019-0020
Rose Elementary (710 W. Michigan St.)	32.17	-110.98	2299	O <sub>3</sub> , PM <sub>10</sub>	PDEQ	SPM	Urban	Population	16670	04-019-1032

<i>Site Index - Ambient Air Monitoring Locations in Arizona in 2006</i>										
<b>City/Site and Address</b>	<b>Lat.</b>	<b>Long.</b>	<b>Elev. (feet)</b>	<b>Parameters Measured</b>	<b>Operator</b>	<b>Classification</b>	<b>Scale</b>	<b>Objective</b>	<b>AAAD ID Number</b>	<b>AQS ID Number</b>
Saguaro NP East (3905 S. Old Spanish Trail)	32.17	-110.69	3080	O <sub>3</sub> , IMPROVE	PDEQ, NPS	SPM, Class I	Urban	Visibility	16474	04-019-0021
Saguaro NP West	32.25	-110.19	2621	Bscat, MET, IMPROVE	ADEQ, NPS	Class I	Regional	Visibility	16475	N/A
Santa Clara (6910 S. Santa Clara Ave.)	32.13	-110.98	2539	PM <sub>10</sub>	PDEQ	SLAMS	Neighborhood	Population	16569	04-019-1026
South Tucson (1601 S. 6th Ave.)	32.2	-110.97	2440	PM <sub>10</sub>	PDEQ	SLAMS (PDEQ)	Neighborhood	Population	16635	04-019-1001
Tangerine (12101 N. Camino De Oeste)	32.43	-111.07	2637	O <sub>3</sub> , PM <sub>10</sub>	PDEQ	SLAMS	Urban	Population	16669	04-019-1018
Tucson Downtown (190 W. Pennington)	32.22	-110.98	2365	CO, O <sub>3</sub>	PDEQ	SLAMS	Neighborhood	Population	16671	04-019-0002
Tucson Fairgrounds (11330 S. Houghton)	32.04	-110.77	3077	O <sub>3</sub>	PDEQ	SLAMS	Neighborhood	Population	16672	04-019-1020

<b>Site Index - Ambient Air Monitoring Locations in Arizona in 2006</b>										
<b>City/Site and Address</b>	<b>Lat.</b>	<b>Long.</b>	<b>Elev. (feet)</b>	<b>Parameters Measured</b>	<b>Operator</b>	<b>Classification</b>	<b>Scale</b>	<b>Objective</b>	<b>AAAD ID Number</b>	<b>AQS ID Number</b>
Tucson Transmissometer Transmitter (U of A Clinical Sci. Bldg 1501 N. Campbell)	32.24	-110.95	2578	Bext	PDEQ, ADEQ	SPM (Urban Haze)	Urban	Urban Haze	16655	N/A
Tucson Transmissometer Receiver (150 W. Congress)	32.22	-110.97	0	Bext	PDEQ, ADEQ	SPM (Urban Haze)	Urban	Urban Haze	16826	N/A
Tucson - U of A Central (1100 N. Fremont Ave.)	32.24	-110.95	2578	Bscat	ADEQ	SPM (Urban Haze)	Neighborhood	Population	16662	04-019-1027
<b>Pinal County</b>										
Apache Junction Fire Station (3955 E. Superstition Blvd. TE)	33.42	-111.50	1748	PM <sub>2.5</sub> , PM <sub>10</sub>	PCAQCD	SLAMS	Neighborhood	Population	16358	04-021-3002
Apache Junction Maintenance Yard (305 E. Superstition)	33.42	-111.54	1750	O <sub>3</sub> , MET	PCAQCD	SLAMS	Neighborhood	Population	16589	04-021-3001

<i>Site Index - Ambient Air Monitoring Locations in Arizona in 2006</i>										
<b>City/Site and Address</b>	<b>Lat.</b>	<b>Long.</b>	<b>Elev. (feet)</b>	<b>Parameters Measured</b>	<b>Operator</b>	<b>Classification</b>	<b>Scale</b>	<b>Objective</b>	<b>AAAD ID Number</b>	<b>AQS ID Number</b>
ASARCO - Hayden Junction (Hwy 177)	33.01	-110.81	1925	SO <sub>2</sub>	ASARCO	SPM	Unknown	Source Impact	16592	N/A
Casa Grande Airport 660 W. Aero Dr.	32.95	-111.76	1410	O <sub>3</sub> , MET	PCAQCD	SLAMS	Neighborhood	Population/ Transport	16367	04-021-3003
Casa Grande Downtown (401 Marshall Rd.)	32.88	-111.75	1378	PM <sub>10</sub> , PM <sub>2.5</sub>	PCAQCD	SLAMS	Neighborhood	Population	16588	04-021-0001
Combs (301 E. Combs Rd. )	33.22	-111.56	1178	O <sub>3</sub>	PCAQCD	SPM	Neighborhood	Population	16657	04-021-3009
Coolidge Maintenance Yard (212 E. Broadway)	32.98	-111.51	1460	PM <sub>10</sub>	PCAQCD	SLAMS	Neighborhood	Population	7446	04-021-3004
Cowtown Road (37580 W. Maricopa)	33.01	-111.99	1214	MET, PM <sub>10</sub>	PCAQCD	SPM	Neighborhood	Population	19347	N/A
Eloy City Complex (620 N. Main St.)	32.76	-111.55	1548	PM <sub>10</sub>	PCAQCD	SLAMS	Neighborhood	Population	16594	04-021-3005

<i>Site Index - Ambient Air Monitoring Locations in Arizona in 2006</i>										
City/Site and Address	Lat.	Long.	Elev. (feet)	Parameters Measured	Operator	Classification	Scale	Objective	AAAD ID Number	AQS ID Number
Mammoth County Complex (118 S. Catalina)	32.72	-110.64	2919	PM <sub>10</sub>	PCAQCD	SLAMS	Neighborhood	Population/Background	16600	04-021-3006
Maricopa (44625 W. Garvey Rd.)	33.05	-112.05	1178	O <sub>3</sub>	PCAQCD	SPM	Neighborhood	Population/Exposure	16656	04-021-3010
Pinal Air Park (Water Well # 2, Marana)	32.51	-111.31	1906	PM <sub>10</sub>	PCAQCD	SLAMS	Regional	Background/Transport	16552	04-021-3007
Pinal County Housing Complex (970 N Eleven Mile Corner Rd.)	32.89	-111.57	1443	MET, PM <sub>10</sub>	PCAQCD	SPM	Microscale	Source Impact	18079	04-021-3011
Queen Valley (10 S. Queen Anne Dr.)	32.29	-111.29	2080	Bscat, IMPROVE, VOC, NO <sub>TL</sub> , O <sub>3</sub>	PCAQCD, ADEQ	SPM (NO <sub>TL</sub> , O <sub>3</sub> ) PAMS (VOC), Class I	Regional	Visibility	16394	04-021-8001
Riverside Maintenance Yard (56964 E. Florence)	33.11	-110.97	1771	PM <sub>10</sub>	PCAQCD	SPM	Neighborhood	Source Impact	21429	04-021-3012

<b>Site Index - Ambient Air Monitoring Locations in Arizona in 2006</b>										
<b>City/Site and Address</b>	<b>Lat.</b>	<b>Long.</b>	<b>Elev. (feet)</b>	<b>Parameters Measured</b>	<b>Operator</b>	<b>Classification</b>	<b>Scale</b>	<b>Objective</b>	<b>AAAD ID Number</b>	<b>AQS ID Number</b>
San Manuel (1st & Douglas Ave.)	32.6	-110.63	1089	SO <sub>2</sub>	ADEQ	SPM	Neighborhood	Source Impact	16397	04-021-2001
Stanfield (36697 W. Papago Dr.)	32.88	-111.96	1296	PM <sub>10</sub>	PCAQCD	SPM	Neighborhood	Population	16636	04-021-3008
<b>Santa Cruz</b>										
Nogales Post Office (300 N. Morley Ave.)	31.34	-110.94	3857	PM <sub>10</sub> , PM <sub>2.5</sub> , MET	ADEQ	SLAMS	Neighborhood	Population	16511	04-023-0004
<b>Yavapai County</b>										
Clarkdale - NW (#2) (northwest of cement plant)	34.78	-112.09	4198	PM <sub>10</sub>	PCC	SPM	Unknown	Source Impact	16626	N/A
Clarkdale - SE (#1) (southeast of CTI flyash silo)	34.77	-112.07	3598	PM <sub>10</sub>	PCC	SPM	Unknown	Source Impact	16628	N/A
Hillside (Sheriff's Repeater Station) Closed June 2005	34° 25'	112° 57'	4918	O <sub>3</sub> , IMPROVE	ADEQ	SPM, Class I	Regional	Background/Transport, Visibility	16315	04-025-0005

<b>Site Index - Ambient Air Monitoring Locations in Arizona in 2006</b>										
<b>City/Site and Address</b>	<b>Lat.</b>	<b>Long.</b>	<b>Elev. (feet)</b>	<b>Parameters Measured</b>	<b>Operator</b>	<b>Classification</b>	<b>Scale</b>	<b>Objective</b>	<b>AAAD ID Number</b>	<b>AQS ID Number</b>
Prescott (221 S. Cortez) Closed 3/01/2005	34° 32'	112° 28'	5210	PM <sub>10</sub>	ADEQ	SPM	Neighbor- hood	Population	16528	04-025-2001
Prescott Valley (7601 E. Civic Circle) Opened 3/12/2003	34.59	-112.33	5104	PM <sub>10</sub>	ADEQ	SPM	Neighbor- hood	Population	18392	04-025-2002
<b>Yuma County</b>										
Dome Valley (5110 S. Avenue 18 E) Opened 5/13/2005	32.75	-114.33	180	MET	ADEQ	SPM	N/A	Special Study	19483	N/A
San Luis (767 N. 1st Ave.) Opened 5/13/2005	32.49	-114.78	112	MET	ADEQ	SPM	N/A	Special Study	18250	N/A
Yuma - Courthouse (2440 W. 28 <sup>th</sup> St.)	32.68	-114.65	98	PM <sub>10</sub>	ADEQ	SLAMS	Neighbor- hood	Population	17027	04-027-0004

<b>Site Index - Ambient Air Monitoring Locations in Arizona in 2006</b>										
<b>City/Site and Address</b>	<b>Lat.</b>	<b>Long.</b>	<b>Elev. (feet)</b>	<b>Parameters Measured</b>	<b>Operator</b>	<b>Classification</b>	<b>Scale</b>	<b>Objective</b>	<b>AAAD ID Number</b>	<b>AQS ID Number</b>
Yuma Game & Fish (9140 E. 28 <sup>th</sup> St.) Opened 4/14/2005	32.68	-114.48	197	O <sub>3</sub>	ADEQ	SLAMS	Neighborhood	Maximum Concentration	18690	04-027-0006
Yuma Mesa (2186 W. County 15th St.) Opened 5/13/2005	32.61	-114.63	190	MET	ADEQ	SPM	N/A	Special Study	19040	N/A
Yuma Supersite (2323 S Arizona Ave)	32.69	-114.61	167	MET, CO, SO <sub>2</sub> , PM <sub>10</sub> , VOCs	ADEQ	SPM	N/A	Special Study	113219	N/A
Yuma Valley (11486 S. Farm Rd.) Opened 5/13/03	32.62	-114.77	89	MET	ADEQ	SPM	N/A	Special Study	19041	N/A
Yuma West	32.74	-114.70	118	MET	ADEQ	SPM	N/A	Special Study	18247	N/A
<b>Mexico</b>										
Agua Prieta Fire Station (Calle 6 & AVE 15)	31.33	-109.55	3936	PM <sub>10</sub> , MET	ADEQ	SPM	Neighborhood	Population	16361	80-026-1000

<i>Site Index - Ambient Air Monitoring Locations in Arizona in 2006</i>										
<b>City/Site and Address</b>	<b>Lat.</b>	<b>Long.</b>	<b>Elev. (feet)</b>	<b>Parameters Measured</b>	<b>Operator</b>	<b>Classification</b>	<b>Scale</b>	<b>Objective</b>	<b>AAAD ID Number</b>	<b>AQS ID Number</b>
Baja	32.57	-115.00	45	MET	ADEQ	SPM	Neighborhood	Population	22242	N/A
Cortez	32.38	-114.87	69	MET	ADEQ	SPM	Neighborhood	Population	22240	N/A
Mexico Supersite	32.47	-114.77	125	MET, CO, PM <sub>10</sub>	ADEQ	SPM	N/A	Special Study	113221	N/A
Sonora Nogales Fire Station (Northwest corner of Lopaz and Mantels)	31.33	-110.94	3943	PM <sub>10</sub>	ADEQ	SPM	Neighborhood	Population	16399	80-026-0005
Sonora	32.42	-114.80	109	MET	ADEQ	SPM	Neighborhood	Population	22243	N/A

*Sites shown in the site index table are based on the best information available at the date of publication.*

N/A – Not available

## Appendix 2 – Acronyms and Abbreviations

ADEQ	Arizona Department of Environmental Quality
ADOT	Arizona Department of Transportation
AgBMP	Agricultural Best Management Practices
APCC	Arizona Portland Cement Co.
APS	Arizona Public Service
Area A	Designated Phoenix metropolitan area
ASARCO	ASARCO LLC - U.S. operating subsidiary of Grupo Mexico
ASU	Arizona State University
$B_{abs}$	Light absorption
$B_{ag}$	Light absorption by gasses
$B_{ap}$	Light absorption by particles
$B_{ext}$	Light extinction
$B_{scat}$	Light scattering
$B_{sg}$	Light scattering by gasses
$B_{sp}$	Light scattering by particles
BACM	Best Available Control Measures
BHP	BHP Copper, Inc.
CAAA	1990 Clean Air Act Amendments
CASTNET	Clean Air Status and Trends Network
CFR	Code of Federal Regulations
Class I	Federally designated park or wilderness area with mandated visibility protection
CMSA	Consolidated Metropolitan Statistical Area
CO	Carbon monoxide
CTOC	Cap and Trade Oversight Committee
Delta T	Difference between two levels of temperature measurements
EPA	U.S. Environmental Protection Agency
FMIC	Ft. McDowell Indian Community
FRM	Federal Reference Method
GRIC	Gila River Indian Community
HAPs	Hazardous Air Pollutants
HART	Hazardous Air Response Team
HC	Hydrocarbon
IMPROVE	Interagency Monitoring of Protected Visual Environments
ITEP	Institute for Tribal Environmental Professionals
km	Kilometers
m	Meters
MAG	Maricopa Association of Governments
MCAQD	Maricopa County Air Quality Division
MET	Meteorological measurements (wind, temperature, relative humidity)
mm	Millimeter
$Mm^{-1}$	Inverse megameter
MSA	Metropolitan Statistical Area

$\mu\text{g}/\text{m}^3$	Micrograms per cubic meter
MSM	Most Stringent Measures
NAAQS	National Ambient Air Quality Standards
NAMS	National Air Monitoring Station
NEAP	Natural Event Action Plan
NM	National Monument
NO	Nitric Oxide
$\text{NO}_2$	Nitrogen Dioxide
$\text{NO}_x$	Sum of NO and $\text{NO}_2$
NPS	National Park Service
$\text{O}_3$	Ozone
PAMS	Photochemical Assessment Monitoring Station
Pb	Lead
PCC	Phoenix Cement Company
PDEQ	Pima County Department of Environmental Quality
PDMI	Phelps Dodge Miami Inc.
PCAQCD	Pinal County Air Quality Control District
PM	Particulate Matter
$\text{PM}_{2.5}$	Particulate Matter $\leq 2.5$ microns
$\text{PM}_{10}$	Particulate Matter $\leq 10$ microns
ppb	parts per billion
ppm	parts per million
Pressure	Barometric air pressure
RH	Relative Humidity
SCE	Southern California Edison
SIP	State Implementation Plan
SLAMS	State and Local Air Monitoring Station
$\text{SO}_2$	Sulfur Dioxide
$\text{SO}_4^-$	Sulfate
SPM	Special Purpose Monitor
SRP	Salt River Project
SRPMIC	Salt River Pima-Maricopa Indian Community
STN	Speciation Trends Network
TEOM	Tapered Element Oscillating Microbalance
TEP	Tucson Electric Power
TSP	Total Suspended Particulates
U of A	University of Arizona
USFS	U.S. Forest Service
VOC	Volatile Organic Compounds
VIOC	Visibility Index Oversight Committee
Wind	Wind speed and direction
WMAT	White Mountain Apache Tribe

## Appendix 3 – Related Web Sites

---

**Air Explorer** (<http://www.epa.gov/airexplorer/>) Air Explorer is a collection of user-friendly visualization tools for air quality analysts. It is linked directly to the EPA's Air Quality Subsystem database.

**AirWeb: Protecting Air Quality** (<http://www2.nature.nps.gov/air/>)  
Learn about how the National Park Service Air Resources Division and the Fish and Wildlife Service Air Quality Branch strive to preserve, protect, enhance and understand the air quality and other resources of our national parks and refuges.

**Arizona Department of Environmental Quality** ([www.azdeq.gov](http://www.azdeq.gov))  
ADEQ's Web site contains information on air quality, news releases, public meetings and many other services that can be provided that help to protect a safe and healthy environment.

**Earth 911: Making Every Day Earth Day!** ([www.earth911.org](http://www.earth911.org))  
That's their mission "to make every day an earth day!" so you can act on today's environmental issues, in order to preserve and maintain for today and tomorrow.

**Earth's Biggest Environment Search Engine** ([www.webdirectory.com](http://www.webdirectory.com))  
This Web site is a directory to numerous environmental subjects, from air to wildlife.

**Environmental Protection Agency** ([www.epa.gov](http://www.epa.gov))  
On EPA's Web site, you can find information about the federal government's role in environmental protection.

**EPA – Air and Radiation** ([www.epa.gov/oar/oaqps](http://www.epa.gov/oar/oaqps))  
You'll breathe easier when you see EPA's air quality planning and standards Web site. They have from what's new in air to the latest projects, programs and contracts.

**EPA's – AIRNow** ([airnow.gov/](http://airnow.gov/))  
Easy access to local air quality forecasts, real-time data, air quality index (AQI), animated color contours of measured AQI values for geographic areas and more.

**EPA's Air Quality Database** ([www.epa.gov/air/data/index.html](http://www.epa.gov/air/data/index.html))  
EPA's air quality database contains extensive air data. On this site, you can find the sources that contribute to emissions, the equipment and facilities that monitor the air, maps on air-related information, and contact information for experts on specific issues regarding air and environment.

**EPA – Region 9** (<http://www.epa.gov/region09/>)

Learn about EPA activities in Arizona, California, Hawaii, Nevada and the Pacific Islands at the Region 9 website.

**FirstGov** ([www.firstgov.gov](http://www.firstgov.gov))

Through this Web site, you can find more than 1,000 federal and state environmental agencies with details about the environment.

**The Interagency Monitoring of Protected Visual Environments Project**

(<http://vista.cira.colostate.edu/improve/>)

On this site, you can take a look at photos of what haze (pollution) can do to the beautiful views of our nation. You can also take a look at what is being done and how you can get involved to improve the views of our nation.

**Inter Tribal Council of Arizona, Inc.** ([www.itcaonline.com](http://www.itcaonline.com))

The site lists the member tribes and includes information about environmental monitoring programs.

**Maricopa County Air Quality Information** ([www.maricopa.gov/envsvc/airqual.asp](http://www.maricopa.gov/envsvc/airqual.asp))

Maricopa County's Environmental Services' Web site has specific descriptions plus current and historical data on the county's air monitors.

**National Tribal Environmental Council** ([www.ntec.org](http://www.ntec.org))

NTEC is a tribal government membership organization with 160 member tribes that work to protect and preserve the reservation environment.

**National Weather Service** ([www.nws.noaa.gov](http://www.nws.noaa.gov))

Dive into the latest occurrences and studies of your weather and atmosphere. There are links to local weather service agencies in each state.

**Visibility Web Cameras** (<http://www.phoenixvis.net>)

This page provides an overview of all Phoenix Visibility Web Cameras. Digital images from Web-based cameras are updated every 15 minutes.

**Pima County Air Quality Information** ([www.deq.co.pima.az.us](http://www.deq.co.pima.az.us))

The Pima County Department of Environmental Quality's Web site has information about air, water and waste programs, and the latest news and regulations that affect Pima County.

**Pinal County Air Quality Information** (<http://co.pinal.az.us/airqual/monitoring.asp>)

Current air quality information from the Pinal County Air Quality Control District.

**Pollen Information** ([www.pollen.com](http://www.pollen.com))

Does it feel like something is in the air? Visit [pollen.com](http://pollen.com) to find out about what kinds of allergens are in your air and when they are there.

**The United States National Park Service** ([www.nps.gov](http://www.nps.gov))

Information about our national parks.

**Visibility Information Exchange Web System (VIEWS)**

(<http://vista.cira.colostate.edu/views/>)

The Visibility Information Exchange Web System is an online exchange of visibility data, research, and ideas designed to support the Regional Haze Rule enacted by the U.S. Environmental Protection Agency (EPA) to reduce regional haze in national parks and wilderness areas. In addition to this primary goal, VIEWS supports global efforts to better understand the effects of air pollution on visibility and to improve air quality in general.

**Weather and Air Quality in the Southwest** ([www.weathersmith.com](http://www.weathersmith.com))

This site contains weather forecasts and air quality information for Phoenix and Tucson.

**Western States Air Resources Council** ([www.westar.org](http://www.westar.org))

WESTAR is composed of 15 western states that have come together to discuss and exchange information on western regional air quality issues.



## Appendix 4 – Maps

---

This section contains maps displaying monitor locations and location information.

### Ambient Air Monitors

This map shows the location of monitors operated by ADEQ, county agencies, private industry and federal agencies.

### Criteria Pollutant Monitoring (Phoenix and Tucson Metropolitan Areas)

These maps identify the locations of monitors of criteria pollutants in Arizona's two largest metropolitan areas.

### Nonattainment and Attainment Areas

This map identifies the areas in Arizona that are nonattainment for PM<sub>10</sub>, SO<sub>2</sub>, CO and O<sub>3</sub>.

### Ozone Network

This map shows the location of ozone monitors operated by ADEQ, private industry, county agencies, and the National Park Service.

### PM10 Network

The location of PM<sub>10</sub> particulate monitors is shown on this map.

### SO2 Network

This map shows the location of the SO<sub>2</sub> monitors and includes the maintenance and nonattainment areas.


### Visibility Network

Urban and regional haze visibility monitoring sites are shown on this map.



### Nephelometers, Transmissometers, Cameras

This map shows the location of each of these types of monitors that ADEQ operates for the study of urban and regional visibility.

# AMBIENT AIR MONITORS




# CRITERIA POLLUTANT MONITORING



Source: AAAD




# Nonattainment and Attainment Areas



- O3-1hr (see below) \*\*
- O3-8hr Nonattainment
- SO2 Nonattainment
- SO2 Attainment with a Mairt Plan
- CO Nonattainment
- CO Attainment with a Mairt Plan
- PM10 Nonattainment
- PM10 Attainment with a Mairt Plan


\*\*May 2005 Redesignated  
June 2005 Revoked



Source: AAD



# Ozone Network



**O3 MONITOR OPERATOR**


- ADEQ
- NATIONAL PARKS SERVICE
- MARICOPA COUNTY
- PIMA COUNTY
- PINAL COUNTY
- SOURCE OPERATED

0 50 100  
miles

Source: AAAD



# P M 1 0 Network




**PM10 MONITOR OPERATOR**

- ADEQ
- ADEQ AND SOURCE
- MARICOPA COUNTY
- PIMA COUNTY
- PINAL COUNTY
- SOURCE


0 50 100  
miles  
Source: AAAD



# SO<sub>2</sub> Network



# VISIBILITY NETWORK



## Nephelometers, Transmissometers, Cameras

